



Management of Leaf Blight and Node Blight Diseases Indian Barnyard Millet (*Echinochloa frumentacea* (Roxb))

**M. Paramasivan ^a, M. Deivamani ^{b*}, S. Thangeswari ^c,
R. Ramjegathesh ^d, A. Thangam ^e, M. Senthilkumar ^f,
S. Malathi ^g and V.K. Satya ^h**

^a Regional Research Station, Tamil Nadu Agricultural University, Vridhachalam, Cuddalore, Tamil Nadu, India.

^b ICAR-Krishi Vigyan Kendra, Tamil Nadu Agricultural University, Pappapattay, Dharmapuri, Tamil Nadu, India.

^c Department of Plant Pathology, Tamil Nadu Agricultural University, Coimbatore, Tamil Nadu, India.

^d National Pulses Research Centre, Tamil Nadu Agricultural University, Pudhukottai, Tamil Nadu, India.

^e Government Arts and Science College for Women, MU, Alankulam, Tenkasi, Tamil Nadu, India.

^f Horticultural Research Station, Tamil Nadu Agricultural University, Yercaud, Salem, Tamil Nadu, India.

^g TNAU Information and Training Centre, Tamil Nadu, India.

^h Horticulture College and Research Institute for Women, Tamil Nadu Agricultural University, Tiruchirapalli, Tamil Nadu, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/arja/2025/v18i1640>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/128862>

Original Research Article

Received: 05/11/2024

Accepted: 07/01/2025

Published: 11/01/2025

*Corresponding author: E-mail: deivamani.m@tnau.ac.in;

Cite as: Paramasivan, M., M. Deivamani, S. Thangeswari, R. Ramjegathesh, A. Thangam, M. Senthilkumar, S. Malathi, and V.K. Satya. 2025. "Management of Leaf Blight and Node Blight Diseases Indian Barnyard Millet (*Echinochloa frumentacea* (Roxb))". *Asian Research Journal of Agriculture* 18 (1):26-31. <https://doi.org/10.9734/arja/2025/v18i1640>.

ABSTRACT

A study on the management of important diseases of rainfed kuthiraivali was carried out through biological control and fungicides in combination with seed treatment and foliar application methods and different intervals of diseases incidence. The leaf blight pathogen was isolated from the affected leaf; the isolated fungus's dark grayish-colored, fluffy mycelia were observed on the PDA medium. The mycelium was grey to olivaceous green, profusely branched, and septate. Conidiophores were single or in small groups, straight to flexuous, septate, smooth, and dark brown. Measuring 3.5 μm x 0.865 μm (1 μm), The disease incidence was observed at 15-day intervals using the percent disease index. Among three years, the seed treatment (10gm/kg) of *Pseudomonas fluorescens* (TNAU Pf1) followed by foliar application of Carbendazim + Mancozeb 2 gm/lit decreases the leaf blight diseases (63.63% reduction over control) and also recorded an increased yield of 1075 kg/ha of kuthiraivali. Seed treatment with *Pseudomonas fluorescens* Pf1 combined with foliar fungicide sprays significantly enhances yield and cost-benefit ratio under rainfed conditions while effectively managing seed-borne and foliar diseases.

Keywords: Leaf blight; seed treatment; foliar fungicide; *Pseudomonas fluorescens*.

1. INTRODUCTION

Barnyard millet (*Echinochloa frumentacea*) is a highly nutritious crop and fast growing robust crop. It is grown even in higher altitudes up to a height of 2000 m in the Himalayas. Barnyard millet serves as dual purpose for feed and fodder crop. It is mainly grown in India, China, Japan, and Korea (Nehru et al., 2023; Govanakoppa et al., 2024). It is cultivated as a substitute for rice on marginal lands where rice and other crops will not grow well (Gomashe, 2017; Mageshwaran et al. 2023; Harb et al., 2024). The grains of barnyard millet are particularly recommended for diabetics and also are appropriate food for patients intolerant to gluten causing celiac disease and have an excellent storage life and can be stored even up to 100 years. The barnyard millet contains 10.5% protein, 3.6% fat, 68.8% carbohydrate and 398 kcal/100 g energy. The total dietary fiber content was high (12.6%) including soluble (4.2%) and insoluble (8.4%) fractions (Kumar et al., 2020; Rawat et al., 2018; Ramesh et al., 2024). Although this crop is hardy, the crop is challenged by many fungal diseases like leaf spot (*Exserohilum monoceras*), blast (*Pyricularia grisea*), grain smut (*Ustilago panici-frumentacei* Bref.) and head smut (*Ustilago crusgalli*) (Prabu et al., 2020). Among these diseases, leaf spot is one of the most destructive diseases after smut, which if occurs at early stages as blight, reduces the crop yield drastically. Larger cultivation of kuthiraivali in climate changes the *Helminthosporium* (*Exserohilum monoceras*) leaf blight incidence was high (47.80%), Leaf spot or blight is incited by the fungus *Exserohilum monoceras*

(Drechsler) Leonard & Suggs [Syns. *Helminthosporium monoceras* (Drechsler); *Helminthosporium crusgalli* (Nisikado and Miyake); *Bipolaris monoceras* (Drechs.) Shoemaker; *Drechslera monoceras* (Drechs.) Subram and Jain; and *Luttrellia monoceras* (Drechsler) Chochryakov]. Morita et al. (2013) described the perfect stage of the fungus as *Setosphaeria monoceras*. Leonard and Suggs (1974) established the genus *Exserohilum* for *Helminthosporium* species and differentiated from its former group *Drechslera* and *Bipolaris* by its stable strongly protuberant conidial hilum a pot experiment was conducted to study the effect of coir pith compost, bone meal powder and panchagavya on barnyard millet (*Echinochloa frumentacea* (Roxb.). Pot experiment revealed that application of 100% RDF (T2) registered the highest quality characters viz., crude protein, crude fiber and ash contents were 5.50, 4.35 and 4.31 per cent, respectively (Srinivasan and Venkatakrishnan, 2021).

2. MATERIALS AND METHODS

2.1 Isolation of Pathogens

Barnyard millet leaves showing typical symptoms of brown spot were collected from the field. Leaves with symptoms were first washed in tap water and then cut into small bits of 2 mm size, containing the discolored brown spots. These bits were surface sterilized with 0.1 per cent mercuric chloride solution for two minutes followed by three changes of sterilized distilled water. These bits were placed on potato dextrose agar under aseptic condition. Inoculated plates

were incubated at 27 °C and watched for any contamination for seven days. After seven days of incubation, fungal colonies completely covered the plates and become dark greyish in color indicating the production of spores.

2.2 Proving Pathogenicity

Pathogenicity test was carried out to prove the Koch's postulates on barnyard millet variety CO (KV2). The conidial suspension of *E. monoceras* at 10^5 spores per ml was inoculated at three to four leaf stage by spray inoculation. The initiation of leaf spot symptom started five to six days of inoculation. Initially dark brown longitudinal spots were observed. Later, several such spots coalesced and covered the major portion of leaf causing blight symptom. The infected leaves turned grey and dried up. Re-isolation of the pathogen was made from artificially inoculated plants and was found to be that of *E. monoceras*.

2.3 Pot Culture Experiments

The pot culture experiments were conducted with eight treatments for seed treatment and foliar spray, Chemical and biocontrol agents.

2.4 Field Experiments

The experimental trial consisting of 9 treatments with biocontrol agents and fungicides was laid out at Dryland Agricultural Research Station Chettinad; the test crop is Kudiraivali CO (Kv2). Randomized Block Design was adopted with three replications. The plot size for each treatment was 20m². The disease incidence of recorded at 15 days intervals. The yield parameters were recorded and analyzed statistically. All normal agronomic practices were followed. In addition, growth parameters viz., plant height, number of branches. Number of ear heads and ear head length and breadth, and seed yield kg/ha.

3. RESULTS AND DISCUSSION

3.1 Leaf Blight Pathogens

The isolated fungus was identified as *Exserohilum monoceras* based on its cultural and morphological characters. The dark grayish coloured fluffy mycelia of the isolated fungus were observed on PDA medium. The mycelium was grey to olivaceous green, profusely branched and septate. Conidiophores were

single or in small groups, straight to flexuous, septate, smooth, dark brown. Conidia pale to mid olivaceous brown, fusoid, smooth, straight or curved, mostly 4-7. Measuring 3, 5 µm x 0.865µm (1 µm).

3.2 Biocontrol Agents and Chemicals against Leaf Blight Diseases of Kudiraivali Under Rainfed Conditions

The experiment result revealed that, leaf blight disease incidence ranged from 15.3 to 40.66 per cent disease incidence (Table 1). Seed treatment *Pseudomonas fluorescens* (10gm/kg of seed) + Foliar spraying of Carbendazim + Mancozeb (2gm/lit) and Seed treatment (10gm/kg of seed) + Seed treatment *Pseudomonas fluorescens* (10gm/kg of seed) + Foliar spraying *Pseudomonas fluorescens* (1gm/lit) less disease incidence recorded, 15.99 and 16.97 on par with each other. The treatment Seed treatment *Pseudomonas fluorescens* (10gm/kg of seed) + Foliar spraying of Carbendazim+ Mancozeb (2gm/lit) for maximum control of leaf blight diseases of Kudiraivali significant reduction for 63.63 and 61.40 for per cent disease reduction comparing with other chemical.

3.3 Yield Analysis

Among the treatments, Seed treatment *Pseudomonas fluorescens* Pf1 (10gm) + Foliar Spray of Carbendazim+ Mancozeb 2gm/lit received for highest yield 1075 kg / ha and increasing cost benefit ratio (1:2.50) under rainfed conditions comparing to other treatments (Table 2).

The similar study reported by Laxmi Rawat et al, (2018) brown spot disease control (75.99) was recorded in the treatment T4 (Soil application of value added vermi-compost impregnated by fluorescent *Pseudomonas* isolate UUHF Psf- 4 + Seed treatment with fluorescent *Pseudomonas* isolate UUHF Psf- 4 + One foliar spray with Fluorescent *Pseudomonas* isolate UUHF Psf- 4 at the time of 50% flowering). Maximum grain yield (23.13 q/ha), highest per cent increase in grain yield (64.86%) and highest per cent avoidable loss (39.34%) Since, the primary infection comes from the seed-borne inoculum; seeds treatment with systemic fungicides before sowing helps in disease control. Spraying of copper fungicides @ 0.3 % helps in reducing the disease intensity (Nagaraja et al., 2007). The management of leaf blight pathogen with

Table 1. Effect of biocontrol agents and chemicals against foliar disease of kudiraivali (Pooled analysis)

S. No	Treatments	Leaf blight PDI [Pooled 2020 & 2021]	Percent reduction over control
T1	ST (10gm) +FS <i>Pseudomonas fluorescens</i> (Pf1) 1gm/lit	16.97 (24.31)	61.40
T2	ST (10gm) +FS <i>Bacillus subtilis</i> (Bs) 1gm/lit	24.15 (29.40)	45.07
T3	ST Pf1 (10gm) +FS-Carbendazim+ Mancozeb 2gm/lit	15.99 (23.55)	63.63
T4	ST (10gm) Bs+ FS Carbendazim+ Mancozeb 2gm/lit	21.24 (27.36)	51.69
T5	FS Carbendazim + Mancozeb 2gm/lit	20.98 (27.24)	52.28
T6	FS Mancozeb 2gm/lit	25.33 (30.26)	42.39
T7	FS Copper oxychloride 2gm/lit	24.30 (29.52)	44.73
T8	FS Carbendazim 1gm/lit	24.40 (29.52)	44.50
T9	Control	43.97 (41.50)	-
	C.D.	3.59	
	SE(d)	1.533	
	C.V.	5.25	

Table 2. Effect of biocontrol agents and chemicals yield on kudiraivali (Pooled analysis)

S. No	Treatments	Yield Kg/ha [Pooled 2020 & 2021]	BCR ratio
T1	ST (10gm) +FS <i>Pseudomonas fluorescens</i> (Pf1) 1gm/lit	1027.50	2.05
T2	ST (10gm) +FS <i>Bacillus subtilis</i> (Bs) 1gm/lit	982.50	2.00
T3	ST Pf1 (10gm) +FS-Carbendazim+ Mancozeb 2gm/lit	1075.00	2.50
T4	ST (10gm) Bs+ FS Carbendazim+ Mancozeb 2gm/lit	1027.50	2.15
T5	FS Carbendazim+ Mancozeb 2gm/lit	997.50	2.08
T6	FS Mancozeb 2gm/lit	925.00	1.97
T7	FS Copper oxychloride 2gm/lit	860.00	1.85
T8	FS Carbendazim 1gm/lit	862.50	1.79
T9	Control	737.50	1.60
	C.D.	85.58	
	SE(d)	36.54	
	C.V.	3.87	

nineteen plant extracts tested under *in vitro* conditions. Among twenty plant extracts the 5v (Vilvum, Vembu, Vanni, Vagai and Vengai) plants extracts (10%) effectively inhibited mycelium growth (84.81, 64.77, 63.26, 61.36 and 48.56 percent reduction over control) of Leaf blight pathogens (Paramasivan et al., 2024).

4. CONCLUSION

This study focused on the management of important diseases of rainfed kudiraivali, which was carried through biological control and fungicides individual and combination with seed treatment and foliar application methods and different intervals of diseases incidence taken the

leaf blight pathogen was isolated from affected leaf, the dark grayish coloured fluffy mycelia of the isolated fungus were observed on PDA medium. Among the treatments, Seed treatment *Pseudomonas fluorescens* Pf1 (10gm) with Foliar Spray of Carbendazim and Mancozeb (2gm/lit) received for highest yield and increasing cost benefit ratio under rainfed conditions comparing to other treatments.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image

generators have been used during writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Gomashe, S. S. (2017). Barnyard millet: present status and future thrust areas. *Millet and sorghum: biology and genetic improvement*, 184-198.
- Govanakoppa, N., B. R., J., S., A., & K. N., J. H. (2024). Recombinant Breeding in Barnyard Millet (*Echinochloa frumentacea* L.): Pioneering High-yield Varieties for Sustainable Agriculture. *Journal of Scientific Research and Reports*, 30(11), 826–833.
<https://doi.org/10.9734/jsrr/2024/v30i112610>
- Harb, H. E., El-Tabakh, M. A., Khattab, A. M., Mohamed, Y. A., Saleh, A. M., & El-Abeid, S. E. (2024). Recent Advances of Using Innovative Strategies in Management of Millet Plant Pathogens. *Genetic improvement of Small Millets*, 297-328.
- Kumar, A., Paliwal, A., Shikha, ., Shyam, R., & Singh, S. B. (2020). Barnyard Millet (*Echinochloa frumentacea*) Varieties Performance under Different Fertility Levels in Rainfed Conditions of Garhwal Himalaya, India. *Current Journal of Applied Science and Technology*, 39(46), 102–106.
<https://doi.org/10.9734/cjast/2020/v39i4631180>
- Laxmi Rawat, T.S. Bisht, Shambhoo Prasad, T. Samuel and Patro, S.K. 2018. Management of Important Endemic Diseases of Barnyard Millet (*Echinochloa frumentacea* L.) by the Use of BioControl Agents in Mid Hills of Uttarakhand, India. *Int.J.Curr.Microbiol.App.Sci.* 7(02): 64-70
- Leonard, K. J. and Suggs, E. G., 1974, *Setosphaeria prolata*, the ascigerous state of *Exserohilum prolatum*. *Mycologia*, 66: 281-297.
- Mageshwaran, V., Paulraj, S., & Nagaraju, Y. (2023). Current insights into the role of rhizosphere bacteria in disease suppression in millets. In *Millet Rhizosphere* (pp. 121-147). Singapore: Springer Nature Singapore.
- Morita, A., Yoshimoto, S., Kosuke, I. and Chihiro, T., 2013, Teleomorph formation of *Setosphaeria monoceras*, a perfect state of *Exserohilum monoceras*, by Japanese isolates. *Mycoscience*, 53(2): 144-146
- Nagaraja, A., Kumar, J., Jain, A. K., Narasimhudu, Y., Raghuchander, T., Kumar, B. and Gowda, H. B., 2007, Compendium of small millets diseases. Project coordination cell. AICSMIP, Univ. Agric. Sci., GKVK, Bangalore, India, pp. 80.
- Nehru, G., Reddy, A. T., Reddy, C. V. C. M., & Sreenivasulu, K. N. (2023). Study of Genetic Diversity in Indian Barnyard Millet Genetic Resources [*Echinochloa frumentacea* (L.)]. *International Journal of Environment and Climate Change*, 13(10), 3094–3110.
<https://doi.org/10.9734/ijecc/2023/v13i102978>
- Paramasivan M, I Johnson, A Thangam, R Ramjegathesh, S Thangeswari and Navarasu. 2024. *Curvularia lunata* causing leaf blight barnyard millet (*Echinochloa frumentacea* (Roxb.) link in India and its management by botanicals. *Int. J. Adv. Biochem. Res* ;8(7S):911-913.
- Prabu, R., Vanniarajan, C., Vetrivanthan, M., Gnanamalar, R. P., Shanmugasundaram, R., & Ramalingam, J. (2020). Diversity and stability studies in barnyard millet (*Echinochloa frumentacea* (Roxb). Link.) germplasm for grain yield and its contributing traits. *Electronic Journal of Plant Breeding*, 11(2), 528-537.
- Ramesh, G. V., Palanna, K. B., Farooqkhan, Rajashekhara, H., Rajesh, F. G., & Das, I. K. (2024). Major Diseases of Small Millets and Their Management Strategies. In *Genetic improvement of Small Millets* (pp. 87-118). Singapore: Springer Nature Singapore.
- Rawat, L., Bisht, T. S., Prasad, S., Samuel, T., & Patro, S. K. (2018). Management of important endemic diseases of barnyard millet (*Echinochloa frumentacea* L.) by the use of bio-control agents in mid hills of Uttarakhand, India. *Int J Curr Microbiol Appl Sci*, 7(2), 64-70.
- Srinivasan, S and D. Venkatakrishnan (2021). Barnyard Millet Quality Characters and NPK Uptake as

Influenced by Integrated Organic
Nutrients in Sandy Loam Soil. Bull.

Env. Pharmacol. Life Sci., Vol 10 [11]: 139-
142.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2025): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/128862>