



Growth and Yield of Sweet Corn as Influenced by Nitrogen Levels and Crop Geometry

Deepak Singh Pachoriya ^{a++}, A.A Kawade ^{b#*}
and V. V. Panchal ^{b#}

^a Agronomy, Shri Vaishnav Institute of Agriculture, SVVV, Indore, India.

^b Shri Vaishnav Institute of Agriculture, SVVV, Indore, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jabb/2025/v28i11883>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/128658>

Original Research Article

Received: 04/11/2024

Accepted: 06/01/2025

Published: 14/01/2025

ABSTRACT

An experiment was conducted during *rabi*, 2023 at new experimental cum demonstration field, Shri Vaishnav Institute of Agriculture, Shri Vaishnav Vidyapeeth Vishwavidyalaya, Indore. The field experiment was laid out in a split plot design with three replication. The main plot treatment comprised of three crop geometry *viz.*, 60 cm x 20 cm (S₁), 60 cm x 30 cm (S₂), 60 cm x 40 cm (S₃) whereas, sub plot treatments comprised of four nitrogen levels *viz.*, 125: 60: 40 kg ha⁻¹ (N₁), 100: 60: 40 kg ha⁻¹ (N₂), 75: 60: 40 kg ha⁻¹ (N₃) and control (N₄). The main objective of research was to study the effect of nitrogen levels and crop geometry on growth and yield of sweet corn. The crop geometry of 60 cm x 20 cm along with application of 125 per cent RDN has shown noticeably

++ M.Sc.;

Assistant Professor of Agronomy;

*Corresponding author: E-mail: kawde.archana@gmail.com;

Cite as: Pachoriya, Deepak Singh, A.A Kawade, and V. V. Panchal. 2025. "Growth and Yield of Sweet Corn As Influenced by Nitrogen Levels and Crop Geometry". *Journal of Advances in Biology & Biotechnology* 28 (1):310-17. <https://doi.org/10.9734/jabb/2025/v28i11883>.

greater growth and yield parameters such as., plant height at harvest, days to 50% flowering, green cob yield, green fodder yield, biological yield and harvest index. While, the crop geometry of 60 cm x 40 cm along with application of 125 per cent RDN has shown noticeably greater growth qualities such as., number of leaves at harvest, leaf area at harvest, dry matter at flowering and at harvest, yield attributing cob length with husk and without husk, cob diameter with husk and without husk, cob weight with husk and without husk, number of grains and grain weight cob⁻¹. Control treatment took maximum days to days to 50 per cent tasselling and 50 per cent silking.

Keywords: Sweet corn; plant spacing; crop geometry; nitrogen fertilizers.

1. INTRODUCTION

Sweet corn, scientifically known as *Zea mays* L. var. *rugosa*, belongs to the Poaceae family, also known as the grass family, encompassing a diverse range of cereal crops such as maize (corn), rice, wheat, barley, and oats. This cultivated variety of maize typically possesses a diploid chromosome number of $2n = 20$, though there can be genetic variation among different cultivars. There is no any other cereal on the earth, that has so immense yield potential as that of maize and hence, occupied a place of "Queen of Cereals".

"Sweet corn (*Zea mays* L. var. *saccharata*) is the same botanical species as common corn, the main difference being that the endosperm in the grains of fresh sweet corn have a greater polysaccharide content at commercial maturity. Sweet corn is prized for its sweet, tender kernels and is primarily grown for human consumption. Sweet corn kernels come in various colours, including yellow, white, and bicolor varieties, with yellow being the most common. However, the sugar in the kernels rapidly converts to starch after its prime harvest stage; sweet corn can be harvested when it reaches the milk stage, while kernel moisture is at approximately 72 to 76 per cent" (Abhishek and Basavanneppa, 2020).

"Sweet corn contains carbohydrates 19 g, sugar 3.2 g, dietary fiber 2.7 g, fat 1.2 g, protein 3.2 g, vitamin a 10 mg, foliate (vit B9) 46 µg, vitamin C 7 mg, iron 0.5 mg, magnesium 37 mg and potassium 270 mg nutritional value per 100 g sweet corn seed. Rich in vitamins, minerals, and dietary fiber, sweet corn provides a healthy addition to the Indian diet. It is particularly valued for its high content of vitamin A, vitamin C, and folate, are essential for vision, immune function and skin, contributing to overall health and well-being. Major sweet corn-producing nations include the United States, China, Brazil, Mexico, Argentina, Thailand, and India. It is cultivated on

nearly 197 m ha with a production of 1148 m tonnes and productivity of 5823.8 kg ha⁻¹ all over the world having wider diversity of soil, climate, biodiversity, and management practices, contributing 37 per cent to global grain production. In the scenario of India, production of sweet corn has been steadily increasing in response to growing domestic demand and expanding export opportunities due to diverse agro-climatic conditions and extensive agricultural infrastructure. The cultivation of sweet corn comes under more than 4 per cent of the net area sown in the country. India produced 30 million tonnes in an area of 9.9 million hectares in 2020-21" (Anonymous 2021). Major sweet corn-producing states include Karnataka, Andhra Pradesh, Maharashtra, Tamil Nadu, and Uttar Pradesh. Madhya Pradesh, located in central India, is renowned for its significant contribution to the country's agricultural sector.

Sweet corn is a heavy feeder, which requires adequate quantity of nitrogen, phosphorus and potassium (Ortas and Sari, 2003) for growth and development. Nutrient requirement of sweet corn varies depending on inherent soil fertility status, cropping season, variety used and management practices. "Besides other factors, the productivity of sweet corn is very much influenced by soil fertility status and quantity of applied nutrient specially nitrogen fertilizers. This nutrient is considered as major yield determining factor required for maize production. But, excessive nitrogen usage can lead to environmental pollution while deficiency results in low plant growth which reduces the grain yield, leaf area duration and rate of photosynthesis. Thus, balanced and optimum use of nitrogen plays a pivotal role in increasing the yield of maize" (Abhishek and Basavanneppa, 2020).

Plant spacing is another important factor which plays a significant role on growth, development and yield of maize. Optimal crop geometry can affect the interception of solar radiation, air

circulation, and nutrient uptake efficiency, thus influencing crop growth and yield. "Maximum yield can be expected only when plant population allows individual plant to achieve their maximum inherent potential"

2. MATERIALS AND METHODS

The experiment was carried out at new experimental cum demonstration field, Shri Vaishnav Institute of Agriculture, Shri Vaishnav Vidyapeeth Vishwavidyalaya, Indore during *rabi* 2023-24.

Soil of the experimental plot was medium and black in colour with good drainage. The topography of experimental field was uniform and fairly levelled. It was low in available nitrogen (210.11 kg ha⁻¹), medium in available phosphorous (14.53 kg ha⁻¹) and very high in available potassium (455.24 kg ha⁻¹). The soil organic carbon content, pH and EC was 0.49 per cent, 7.32 and 0.74 dSm⁻¹ respectively.

The experiment was laid in split plot design with three main plot treatment of three crop geometry viz., 60 cm x 20 cm (S₁), 60 cm x 30 cm (S₂), 60 cm x 40 cm (S₃) whereas, sub plot treatments of four nitrogen levels viz., 125: 60: 40 kg ha⁻¹ (N₁), 100: 60: 40 kg ha⁻¹ (N₂), 75: 60: 40 kg ha⁻¹ (N₃) and control (N₄). The soil was brought to fine tilth and then the ridges and furrows were opened at 60 cm apart. The field was divided into 36 plots with gross plot size of 3.60 m x 4.80 m each.

The recommended dose of phosphorus and potassium (60:40 P₂O₅ and K₂O kg ha⁻¹) was applied to all plots except control treatment before sowing. Two split dose of nitrogen was given as per treatments to individual plots as basal and top dressing. The nutrients were applied through urea, single super phosphate (SSP) and muriate of potash (MOP).

The seeds of sweet corn variety sweet-80 procured from the local market of Indore were used during experimentation. The seeds of sweet-80 provide long uniform cylindrical cobs, good plant vigorous, golden yellow kernels. The seeds were shown at 3-4 cm depth and different spacings according to treatment details. Sowing was done through hand dibbling method. One seeds were sown at each hill.

The various biometric observations were recorded on five randomly selected sweet corn plants. Bamboo pegs along with tags were fixed

at the north side of each plant for easy identification. The growth observations were recorded on these plants starting from 30 DAS at an interval of 15 days up to sweet corn harvesting. These plants were further used for recording yield contributing characters.

The cobs were harvested at milky stage. Harvesting was usually done in the morning when moisture per cent was high and the temperature was low. After the last picking, green fodder from net plot was harvested and weighed for individual plot and final green fodder yield was expressed in kg ha⁻¹.

The statistical analysis of the data was carried out by the standard statistical method 'Analysis of Variance' (Panse and Sukhatme, 1967). The appropriate standard error (SE) and critical difference (CD) at 5 per cent level of significance were worked out.

3. RESULTS AND DISCUSSION

3.1 Effect of Crop Geometry

3.1.1 Growth attributes

The plant height of sweet corn increased progressively from initial growth phase up to crop maturity. The number of leaves and leaf area increased up to 60 days and decreased thereafter towards maturity due to leaf senescence. Total dry matter increased from flowering to harvest. The silking in crop took plant immediately after 4-5 days of tasselling.

The data presented in Table 1 reveals that, the highest plant height (185.08 cm) at harvest was recorded with crop geometry of 60 cm x 20 cm and also took for days to 50 % tasselling and silking i.e. 58.11 and 63.08, respectively. The lowest plant height (168.09 cm) was recorded in treatment with crop geometry of 60 cm x 40 cm. However, the other growth parameters such as no. of leaves plant⁻¹ (13.74), leaf area plant⁻¹ (30.58), dry matter plant⁻¹ (206.81 g) at harvest was recorded in treatment with application having 60 cm x 40 cm crop geometry. The lower value of the above-mentioned parameters was recorded in plots with crop geometry of 60 x 20 cm.

The increased plant height in higher plant density might be due to thick plant stand. As the intensity of shading increases due to more population, the plant tends to grow taller; such increase in height

of the plant at higher population was reported by Agasibagil et al., (2006), Kole (2010), Babu et al., (2019).

Wider crop geometry produced more leaves and leaf area plant⁻¹ than narrow spacing, which may be due to the efficient utilization of growth resources like sunlight, moisture, space, and

nutrients. The higher dry matter production under wider spacing may be due to more availability of nutrients resulted into increase in cell division, assimilation rate and metabolic activities in plant. These findings are substantiated by the findings of Agasibagil et al., (2006), Kole (2010), Babu et al., (2019), Saha et al., (2023) and Ibeawuchi et al., (2008).

Table 1. Growth attributing characters of sweet corn as influenced by nitrogen levels and crop geometry

Treatment	Growth attributes					
	Plant height (cm)	Number of leaves plant ⁻¹	Leaf area plant ⁻¹ (dm ²)	Dry matter plant ⁻¹ (g)	Days to 50 % Tasselling	Days to 50 % Silking
Crop geometry						
S ₁ : 60 cm x 20 cm	185.08	12.47	26.33	190.34	58.11	63.08
S ₂ : 60 cm x 30 cm	182.88	13.24	28.70	202.09	57.88	62.83
S ₃ : 60 cm x 40 cm	176.09	13.74	30.58	206.81	56.29	61.31
S. Em. ±	1.13	0.19	0.66	2.24	0.15	0.16
CD at 5%	3.31	0.56	1.92	6.52	0.43	0.47
Nitrogen levels						
N ₁ : 125 % RDN	187.73	14.01	30.21	212.47	53.93	58.87
N ₂ : 100 % RDN	184.59	13.24	28.89	206.91	56.60	61.62
N ₃ : 75 % RDN	180.76	12.88	27.60	195.38	59.38	64.36
N ₄ : Control	172.32	12.47	27.46	184.23	59.79	64.79
S. Em. ±	1.27	0.28	0.48	1.91	0.28	0.28
CD at 5%	3.70	0.82	1.40	5.58	0.83	0.81
Interaction effect						
A x B	NS	NS	NS	NS	NS	NS
General mean	181.35	13.15	28.54	199.75	57.43	62.41

Table 2.a. Yield attributing characters of sweet corn as influenced by nitrogen levels and crop geometry

Treatment	Yield attributing characters				
	Cobs plant ⁻¹	Cob length (with husk)	(without husk)	Cob Diameter (with husk)	(without husk)
A) Crop geometry					
S ₁ : 60 cm x 20 cm	1.29	18.75	14.69	5.04	4.29
S ₂ : 60 cm x 30 cm	1.30	19.96	16.09	6.58	5.85
S ₃ : 60 cm x 40 cm	1.27	21.25	17.33	7.04	6.33
S. Em. ±	0.02	0.35	0.48	0.17	0.17
CD at 5%	NS	1.02	1.41	0.48	0.50
B) Nitrogen level					
N ₁ : 125 % RDN	1.32	21.68	17.66	7.10	6.33
N ₂ : 100 % RDN	1.29	21.04	17.04	6.61	5.90
N ₃ : 75 % RDN	1.27	19.77	15.72	5.80	5.10
N ₄ : Control	1.25	17.46	13.73	5.37	4.63
S. Em. ±	0.03	0.23	0.30	0.20	0.19
CD at 5%	NS	0.68	0.88	0.57	0.54
C) Interaction effect					
A x B	NS	NS	NS	NS	NS
General mean	1.28	19.99	16.04	6.22	5.49

Table 2. b. Yield attributing characters of sweet corn as influenced by nitrogen levels and crop geometry

Treatment	Yield attributing characters				
	Cob weight (plant ⁻¹)		No. of grains cob ⁻¹	No. of grain rows cob ⁻¹	Grain weight cob ⁻¹
(with husk)	(without husk)				
A) Crop geometry					
S ₁ : 60 cm x 20 cm	208.50	186.00	414.96	15.17	155.00
S ₂ : 60 cm x 30 cm	211.00	190.58	426.43	16.57	161.83
S ₃ : 60 cm x 40 cm	213.50	193.50	432.42	17.80	164.92
S. Em. ±	0.88	1.31	2.15	0.46	1.63
CD at 5%	2.57	3.83	6.27	1.35	4.77
B) Nitrogen level					
N ₁ : 125 % RDN	215.56	195.33	441.46	18.09	167.00
N ₂ : 100 % RDN	213.00	192.44	425.93	17.52	165.22
N ₃ : 75 % RDN	210.22	189.11	418.50	16.22	159.22
N ₄ : Control	205.22	183.22	412.51	14.21	150.89
S. Em. ±	1.15	1.62	5.47	0.30	1.18
CD at 5%	3.35	4.72	15.97	0.88	3.43
C) Interaction effect					
A x B	NS	NS	NS	NS	NS
General mean	211.00	190.03	424.60	16.51	160.58

3.1.2 Yield attributes

As per the data presented in Tables 2.a and 2.b indicates that, the yield attributing characters viz., length of cob with (21.25 cm) and without husk (17.33 cm), diameter of cob with (7.04) and without husk (6.33 cm), fresh weight of cob with (213.50 g) and without husk (193.50 g), no. of grains cob⁻¹ (432.42), no. of grain rows cob⁻¹ (17.80) and grain weight cob⁻¹ (164.92) were found highest in crop geometry of 60 cm x 40 cm spacing. Whereas the minimum values were recorded from the treatment crop geometry of 60 cm x 20 cm spacing. The values of number of cobs plant⁻¹ as influenced by different treatments showed non-significant results.

The yield attribute values decreased as the plant population increased. These results indicate that there is a positive relationship between crop spacing and yield attributing characters probably due to variable plant competition. The enhanced yield under wider planting geometry might be due to significant improvement in overall crop growth of sweet corn by increased photosynthetic efficiency. Earlier researchers, viz., Kar et al., (2006), Mahapatra et al., (2006) and Kanakdurga et al., (2012) reported the same trend regarding

plant geometry and yield attributed of different crops.

3.1.3 Yield

The green cob yield, green fodder yield, biological yield of sweet corn was significantly influenced by different crop geometry. Treatment with crop geometry of 60 cm x 20 cm recorded maximum green cob yield (19157.93 kg ha⁻¹), green fodder yield (35474.94 kg ha⁻¹) and biological yield (54632.87 kg ha⁻¹) which was found to be at par with crop geometry of 60 cm x 30 cm.

Even though the growth and yield parameters were maximum in 60 cm x 40 cm spacing, higher yields under plant spacing of 60 cm x 20 cm was increased yield owing to the maximum number of marketable cobs compared with those of 60 cm x 30 cm and 60 cm x 20 cm spacing. Decrease in crop geometry significantly increased crop cob and green fodder yield resulting in maximum biological yield. These findings are substantiated by the findings Verma and Tomar (2014), Pagar et al., (2022) and saha et al., (2023).

The harvest index of sweet corn as influenced by different treatment was found to be non-significant.

Table 3. Green cob yield, green fodder yield, biological yield and harvest index of sweet corn as influenced by nitrogen levels and crop geometry

Treatment	Yield of sweet corn			
	Green cob yield (kg ha ⁻¹)	Green fodder yield (kg ha ⁻¹)	Biological yield (kg ha ⁻¹)	Harvest index
A) Crop geometry				
S ₁ : 60 cm x 20 cm	19157.93	35474.94	54632.87	34.61
S ₂ : 60 cm x 30 cm	18673.69	34355.92	53029.61	34.98
S ₃ : 60 cm x 40 cm	14903.85	31049.68	45953.53	32.14
S. Em. ±	830.46	861.35	1593.59	0.74
CD at 5%	2424.06	2514.23	4651.59	NS
B) Nitrogen level				
N ₁ : 125 % RDN	21913.27	39297.51	61210.78	35.80
N ₂ : 100 % RDN	18885.74	35088.99	53974.73	34.81
N ₃ : 75 % RDN	16023.33	31919.24	47942.57	32.94
N ₄ : Control	10118.70	28201.66	31269.95	32.08
S. Em. ±	1212.17	1686.14	2834.04	0.80
CD at 5%	3538.23	4921.74	8272.39	2.33
C) Interaction effect				
A x B	NS	NS	NS	NS
General mean	17096.64	33626.85	49716.29	33.91

3.2 Effect of Nitrogen Level

3.2.1 Growth attributes

The plant height of sweet corn increased progressively from initial growth phase up to crop maturity. The number of leaves and leaf area increased up to 60 days and decreased thereafter towards maturity due to leaf senescence. Total dry matter increased from flowering to harvest. The silking in crop took place immediately after 4-5 days of tasselling.

The highest values for growth parameters such as plant height (187.73 cm), no. of leaves plant⁻¹ (14.01) at harvest, leaf area plant⁻¹ (30.21) at harvest, dry matter plant⁻¹ (212.47 g) at harvest was recorded in treatment with application of 125 per cent RDN. The lower value of the above-mentioned parameters was recorded in control treatment. While days required for 50 per cent tasselling (59.79) and silking (64.79) were more in control treatment.

Increase in the fertilizer levels might have increased photosynthate formation and partitioning to stems that might have favourable impacts on plant height of maize. Similar findings in corn were also reported by Singh et al., (2012), Thakur et al., (2015) and Araujo et al., (2005). The readily available nitrogen in soil

and for higher absorption of nutrients must have resulted in increase in number of leaves plant⁻¹. It may be due to increase in assimilation rate, cell division and metabolic activities in plant. Similar results were reported by Tiwari et al., (2021), Shrivankumar et al., (2023). The increase in leaf area with higher nitrogen levels may be because of increased amount of cellular protoplasm. This resulted in the expansion of the cell wall, which was manifested in the increased linear and lateral dimensions of leaves of the plant. Similar results were reported by Thakur et al., (2015) and Sravankumar et al., (2023).

3.2.2 Yield attributes

Yield attributing characters viz., length of cob with (21.68 cm) and without husk (17.66 cm), diameter of cob with (7.10 cm) and without husk (6.33 cm), fresh weight of cob with (215.56 g) and without husk (195.33 g), no. of grains cob⁻¹ (441.46), no. of grain rows cob⁻¹ (18.09) and grain weight cob⁻¹ (167.00) were found highest in treatment with application of 125 per cent RDN. Whereas the minimum values were recorded from control treatment.

The values of number of cobs plant⁻¹ as influenced by different treatments showed non-significant results.

This evidently proved that increased availability of nitrogen to crop at higher levels resulted in production of photosynthates and their efficient translocation for development of reproductive parts. Similar results were reported by Abhishek and basavanneppa (2020) and Thakur et al., (2015).

Thus, greater availability of photosynthates, metabolites and nutrients to develop reproductive structures seems to have resulted in increased productive plants like cob girth, cob length, cob weight and weight of grains of cob with wider spacing and higher nitrogen levels.

3.2.3 Yield

The green cob yield, green fodder yield, biological yield of sweet corn was significantly influenced by different nitrogen levels. Treatment with application of 125 per cent nitrogen recorded maximum green cob yield (21913.27 kg ha⁻¹), green fodder yield (39325.49 kg ha⁻¹) and biological yield (61238.76 kg ha⁻¹) which was found to be at par with application of 100 per cent RDN.

The fresh cob yield in sweet corn was increased with application of 125 per cent RDN over other nitrogen fertilizer levels. The yield of the sweet corn crop is a function of several yield components, which are dependent on complementary interaction between vegetative and reproductive growth of the crop. Increase in growth and yield attributes along with increased nitrogen level must have resulted in maximum green yield of sweet corn. The present findings are in close agreement with the results obtained by Thakur et al., (2015), Singh et al., (2000), Bhadu et al., (2015). and Khan et al., (2018).

The increased fresh cob yield and green fodder yield which ultimately increase biological yield in sweet corn with application of 125 per cent RDN over other treatments. This might be due to the fact that higher levels of nitrogen leads to adequate supply of nutrients to the plant resulting in better growth which in turn lead to better physiological process and movement of photosynthetic to sink. The present findings are in close agreement with the results obtained by Singh et al., (2000)

Increase in nitrogen fertilizer level has resulted to increase in green cob yield and fodder yield. Thus, the harvest index was also found to be increased.

3.2.4 Interaction effect

The interaction effect of crop geometry and nitrogen fertilization in respect of growth attributes, yield attributes and yield was found to be non-significant.

4. CONCLUSION

It can be concluded that sowing of sweet corn with 60 cm x 20 cm along with application of 125 per cent RDN resulted in higher plant height, green cob yield and biological yield which was at par with treatment with 60 cm x 30 cm spacing and 125 per cent RDN application while all other growth and yield parameters were found to be highest in spacing of 60 cm x 40 cm.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Abhishek, N., & Basavanneppa, M. A. (2020). Effect of plant densities and nitrogen levels on cob yield and quality parameters of sweet corn (*Zea mays* L. Saccharata) in irrigated ecosystem. *International Journal of Chemical Studies*, 8(2), 2918–2921.
- Agasibagil, A. B. (2006). Response of maize (*Zea mays* L.) genotypes to planting density in drill sown paddy tract (Master's thesis). University of Agricultural Sciences, Dharwad, India.
- Anonymous. (2021). Maize outlook report— January to May 2021. *Agricultural Market Intelligence Centre*, ANGRU, Lam.
- Araujo, E. D., Feitosa, F. M., Silva, F. C. S., Andrade Junior, I. O. A., Rodrigues, B. R. A., & Mota, W. F. (2017). Growth and yield of baby corn as influenced by nitrogen topdressing. *African Journal of Agricultural Research*, 12(12), 963–969.
- Babu, R., Rampyare, & Kumar, P. (2019). Growth and grain yield of hybrid maize (*Zea mays* L.) under different plant

- geometry and fertility levels. *International Journal of Current Microbiology and Applied Sciences*, 8(11), 450–457.
- Ibeawuchi, I. I., Matthews, E. O., Chinyere, & Onyia, A. (2008). Plant spacing, dry matter accumulation and yield of local and improved maize cultivars. *Journal of American Science*, 4(1), 1545–1003.
- Kanakadurga, S. V., Sreelatha, & Vishnu, V. (2012). Influence of planting methods, spacing and fertilization on yield and quality of sweet corn. *International Journal of Maize Research and Related Industries*, 1(2), 121–123.
- Kar, P. P., Barik, K. C., Mahapatra, P. K., Garnayak, L. M., Rath, B. S., Bastia, D. K., & Khanda, C. M. (2006). Effect of planting geometry and nitrogen on yield, economics and nitrogen uptake of sweet corn (*Zea mays*). *Indian Journal of Agronomy*, 51, 43–45.
- Kole, G. S. (2010). Response of corn (*Zea mays* L.) to plant density and fertilizer level (Master's thesis). University of Agricultural Sciences, Dharwad, India.
- Mahapatra, P. K., Barik, K. C., & Khanda, C. M. (2006). Effect of planting geometry and nitrogen on yield economics and nitrogen uptake ha⁻¹ of corn (*Zea mays* L.). *Indian Journal of Agronomy*, 51(1), 42–45.
- Ortas, I., & Sari, N. (2003). Enhanced yield and nutrient content of sweet corn with mycorrhizal inoculation under field conditions. *Agricultura Mediterranea*, 3(4), 188–195.
- Pagar, P. A., Pawar, S. B., Asewar, B. V., & Patil, D. K. (2022). Effect of spacing and nutrient management practices on growth, yield and economics of sweet corn-chickpea under sequence cropping. *The Pharma Innovation Journal*, 11(7), 280–286.
- Panse, V. G., & Sukhatme, P. V. (1967). *Statistical method for agricultural workers*. Indian Council of Agricultural Research.
- Saha, I., Jat, S. L., Singh, P., Padhan, S. R., Radheshyam, Mandal, A., Ramniwas, & Karkraliya, M. (2023). Influence of planting density and nitrogen management on growth and productivity of maize in eastern India. *Maize Journal*, 12(1), 45–51.
- Singh, U., Saad, A. A., Ram, T., Chand, L., Mir, S. A., & Aga, A. A. (2012). Productivity, economics and nitrogen-use efficiency of sweet corn (*Zea mays saccharata*) as influenced by planting geometry and nitrogen fertilization. *Indian Journal of Agronomy*, 57(1), 43–48.
- Sravankumar, L., Pradeep, R., Kanaujiya, P. K., Gowd, K. A., & Mallikarjun. (2023). Effect of nitrogen and phosphorus levels on growth and yield of maize (*Zea mays* L.). *The Pharma Innovation Journal*, 12(9), 1006–1008.
- Thakur, A. K., Thakur, D. S., Patel, R. K., Pradhan, A., & Kumar, P. (2015). Effect of different plant geometry and nitrogen levels, in relation to growth characters, yield and economics on sweet corn (*Zea mays sachharata* L.) at Bastar plateau zone. *Effect of Different Plant Geometry and Nitrogen Levels*, 10(3), 1223–1226.
- Tiwari, K. D., Chaturvedi, D. P., Singh, T., Yadav, T. K., & Awadhiya, P. (2021). Effect of nitrogen and sulphur levels on growth, yield and quality of maize (*Zea mays* L.). *The Pharma Innovation Journal*, 11(1), 418–422.
- Verma, A., & Tomar, G. S. (2014). Effect of planting geometry and nitrogen levels on growth, green cob yield and economics of sweet corn (*Zea mays saccharata* Sturt.). *International Journal of Agricultural Sciences*, 10, 766–769.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2025): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/128658>