



Comparative Analysis of Ultrasound and Thermography for Detecting Tissue Alterations in Breast Cancer-related Lymphedema

Raiana Fernandes Mariz Simões ^a,
Laura Raynelle Patriota Oliveira ^a,
Ana Cláudia Souza da Silva ^a,
Ana Rafaela Cardozo da Silva ^a,
Maria Gabriela Amaral Lima ^a, Naiany Tenório ^a,
Vanessa Maria da Silva Alves Gomes ^a and Diego Dantas ^{a*}

^a Department of Physical Therapy, Federal University of Pernambuco, Brazil.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jammr/2024/v36i125672>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/128074>

Original Research Article

Received: 11/10/2024

Accepted: 13/12/2024

Published: 17/12/2024

ABSTRACT

Aims: To associate the results of thermography and ultrasound of the upper limb with lymphedema in women after breast cancer.

Study Design: Cross-sectional study.

*Corresponding author: E-mail: diego.sdantas@ufpe.br;

Cite as: Simões, Raiana Fernandes Mariz, Laura Raynelle Patriota Oliveira, Ana Cláudia Souza da Silva, Ana Rafaela Cardozo da Silva, Maria Gabriela Amaral Lima, Naiany Tenório, Vanessa Maria da Silva Alves Gomes, and Diego Dantas. 2024. "Comparative Analysis of Ultrasound and Thermography for Detecting Tissue Alterations in Breast Cancer-Related Lymphedema". *Journal of Advances in Medicine and Medical Research* 36 (12):236-45. <https://doi.org/10.9734/jammr/2024/v36i125672>.

Place and Duration of Study: Conducted in Recife, Brazil, from September 2022 to July 2024.

Methodology: A total of 43 breast cancer survivors were included. Each participant underwent frontal thermography using a Thermovision FLIR Systems C5 camera. Minimum, mean, and maximum temperatures were evaluated at specific points on the forearm (TA) and arm (TB) using the FLIR Tools software. The same points were analyzed using ultrasound to identify fibrosis, fat infiltration, thickening of the dermo epidermal layer (DEC), and loss of differentiation between layers. Statistical analysis was performed using JASP software.

Results: The study identified a higher prevalence of DEC thickening in the arm (68.5%) and forearm (57.9%). However, no significant correlation was observed between skin temperature and tissue alterations ($P > .05$), with a small effect size and limited clinical relevance

Conclusion:

The study revealed a significant prevalence of fibrosis and dermoepidermal layer thickening in lymphedema-affected upper limbs, with higher relative risk observed in participants with lymphedema. Thermographic analysis, however, did not show significant temperature differences correlating with ultrasound findings, highlighting its limited standalone diagnostic value.

Keywords: Lymphedema; breast cancer; ultrasound; thermography.

1. INTRODUCTION

Lymphedema is characterized by the accumulation of protein-rich fluid in the interstitial space due to lymphatic system failure. It is a chronic, progressive, and currently incurable condition (Shen et al., 2023). The prevalence of lymphedema is high among women diagnosed with breast cancer. The main causes of this condition include treatments such as radiotherapy and lymph node removal, which can reduce lymphatic reabsorption and transport capacity. The management of lymphedema is challenging due to diagnostic difficulties and various associated tissue alterations (Levenhagen et al., 2017).

There is no single tool to assess lymphedema; instead, a combination of methods is employed for more accurate conclusions. The most common diagnosis is based on clinical analysis and increased volume of the affected limb. The lack of standardization limits understanding of the disease's incidence and the effectiveness of available treatments (Donahue et al., 2023). An effective tool for diagnosing lymphedema should go beyond simply detecting arm volume increase after surgery. It should be cost-effective, highly accurate, and assist clinicians in understanding the tissue changes associated with the condition. Additionally, it should facilitate the detection of lymphedema in subclinical stages, when arm volume changes are not yet measurable (Shavit et al., 2018).

Lymphedema classification is done in stages, with stage 0 characterized by sensations of heaviness and fatigue in the limb without visible

swelling, which, like stage 1, is reversible (Denlinger et al., 2018). However, diagnostic methods often focus on arm circumference measurement, water displacement, and bioimpedance analysis, comparing the limbs (Yusof et al., 2012). Therefore, imaging and functional methods are becoming increasingly important in research and clinical practice, rather than relying solely on circumference measurements to quantify arm volume increase, despite their simplicity. Thermographic devices, for example, capture infrared radiation emitted by the body and convert it into electrical signals, generating a thermogram that displays temperatures using colors. Although this method is effective in detecting various vascular conditions, skin and tissue problems, and lymphedema, there is still a lack of comprehensive evidence-based insights to support its application in different areas of medicine (Kelly-Hope et al., 2021; Kesztyüs et al., 2023).

While thermography results for diagnosing lymphedema are promising and suggest benefits in identifying patterns associated with tissue changes in lymphedema, such as edema, fibrosis, and liposubstitution (Ibarra Estupiñán et al., 2020), caution is essential, and the accuracy of these documented associations must be evaluated.

Ultrasound, on the other hand, can more accurately detect pathologies in superficial tissues. Studies comparing physical examination with ultrasound imaging reveal that physical examination is ineffective in identifying the more advanced stages of lymphedema, unlike

ultrasound images (Ricci et al., 2022). Changes in stages two and three, considered irreversible, go beyond swelling and include morphological changes in the skin, subcutaneous tissue, and muscles, requiring a trained professional to properly identify them through ultrasound (Goudarzi et al., 2023).

The evaluation of skin and tissue changes by ultrasound is crucial for monitoring and managing women with secondary lymphedema due to breast cancer (Mander et al., 2023). This imaging technique provides a detailed view of the skin layers and underlying tissue, enabling early detection of structural changes such as skin thickening, fibrosis, and fat accumulation (Polat et al., 2020). Identifying these tissue changes allows for a more accurate classification of lymphedema severity. However, no studies to date have related these changes to superficial skin temperature measured by thermography.

This article generally aimed to verify the association between upper limb ultrasound alterations and superficial skin temperature in women with lymphedema, assessed through infrared thermography. Specifically, it aimed to clinically characterize the sample of women with lymphedema; identify qualitative ultrasound alterations (fibrosis, fat infiltration, fluid presence, and loss of subcutaneous tissue differentiation) in the upper limb; and identify the maximum, mean, and minimum superficial skin temperatures of the upper limb through thermography.

2. METHODOLOGY

This was a cross-sectional study analyzing ultrasound and thermographic exams of women with a history of breast cancer, with and without breast cancer-related lymphedema (BCRL). This study is part of a larger research project aimed at evaluating the diagnostic accuracy of thermography in diagnosing BCRL. The research was conducted at the Laboratory of Women's Health and Pelvic Floor Physical Therapy (LAFISMA), Department of Physical Therapy, Federal University of Pernambuco (UFPE), from September 2022 to July 2024. The research protocol was reviewed and approved by the UFPE Research Ethics Committee under protocol no. 5.434.586. All participants provided informed consent by signing the written Informed Consent Form.

For this study, all women with available ultrasound and thermographic exams were included. Eligibility criteria for the primary study included women aged 40 to 70 years with a history of unilateral mastectomy and breast cancer Stage I-II. Exclusion criteria included women with bilateral breast cancer, bilateral mastectomy, primary lymphedema, edema related to other causes (e.g., rheumatologic, renal, neurological, orthopedic problems, or prior vascular disease), skin conditions (erysipelas, intertrigo, or ulcers), and those undergoing chemotherapy or radiotherapy treatment.

BCRL assessment was conducted following the guidelines of the International Society of Lymphology (International Society of Lymphology, 2016). The standard method for lymphedema evaluation consisted of indirect volumetry using the truncated cone volume calculation. This method is characterized by good diagnostic accuracy, excellent reproducibility, and is based on the calculation of the total volume of the affected limb and its comparison with the unaffected limb (Levenhagen et al., 2017).

Data collection was performed at the LAFISMA-UFPE facilities, and evaluators were previously trained and calibrated for all assessments included in the project, which comprised thermographic imaging acquisition, clinical lymphedema examination, and ultrasound evaluation of lymphedema. The use of thermography as an alternative diagnostic method for lymphedema followed the recommendations of the American Academy of Thermology. The thermography process consisted of four key steps: image acquisition, processing, region of interest delimitation, and analysis (Fernández-Cuevas et al., 2017).

Thermograms were obtained in a windowless room at a controlled temperature of 22°C and 60% humidity, regulated by a digital weather station. The room was free from direct sunlight, air drafts, and electrical equipment generating heat. Prior to the examination, participants were instructed to avoid applying creams or perfumes to the skin, consuming stimulants or caffeinated substances, using nasal decongestants, or engaging in vigorous physical exercise within two hours before the exam. Upon arrival, participants remained seated in the room for 20 minutes with exposed upper limbs positioned on their laps to achieve thermal

equilibrium with the room temperature (International Society of Lymphology, 2016).

Superficial skin temperature measurements were obtained using a thermographic camera (Thermovision FLIR Systems C5, resolution 160 x 120, 19,200 pixels). The camera was positioned 1 meter from the participant to capture all regions of interest. Participants were positioned frontally in an anatomical position, with the chest and upper limbs uncovered. Two equidistant points from the elbow joint line were marked: TB (arm point) and TA (forearm point) (Fig. 1).

Thermograms were processed using FLIR Tools software, with a standard emissivity of .98, Rainbow scale, and a temperature range from 23°C to 37.7°C. Each region of interest (ROI) was analyzed using the line measurement tool, with trained evaluators marking 10 cm below and above the cubital crease, referred to as TA and TB, respectively (Polat et al., 2020). For each ROI, the software provided maximum, mean, and minimum temperatures in degrees Celsius (°C). Thermograms were quantitatively interpreted based on the minimum, maximum, and mean temperatures measured, comparing the affected and contralateral limbs, while considering the theoretical framework of thermoregulation (Fernández-Cuevas et al., 2017) and the

pathophysiological process of lymphedema (Cohen, 2009).

Ultrasound images were acquired using GE's LOGIC V5 equipment with a 7–12 MHz linear transducer in B-mode at a depth of 4 cm. The transducer with gel was positioned on the anterior surface of the arm and forearm, following the same orientation as the thermography. The images (Fig. 2) were qualitatively analyzed by a trained evaluator to identify fibrosis, fat infiltration, dermo epidermal layer thickening, and loss of differentiation between the dermis and epidermis at the same points where temperatures were collected (Mander et al., 2019).

Data were stored and processed using JASP software, version 0.18.3.0. Central tendency measures were described using means and standard deviations. Data normality was assessed using the Kolmogorov-Smirnov test. The prevalence ratio for ultrasound findings was calculated using the Odds Ratio (95% CI). The confidence interval (95%) for non-parametric data used to associate thermography with ultrasound was calculated using the Hodges-Lehmann method. Comparisons of skin temperatures according to the ultrasound findings were conducted using the Mann-Whitney test for non-parametric data, and effect sizes were estimated using Hedges' *g*, where values between 0.2 and 0.5 are considered small,




Fig. 1. Schematic thermographic image in the frontal plane, showing the location of the points TA: forearm point and TB: arm point




Fig. 2. Ultrasound images of the upper limb with lymphedema (A and C) and without lymphedema (B and D)

A: affected limb, forearm point; thickening of the dermoepidermal layer and loss of differentiation between the layers can be observed through the blurring effect, as well as fibrosis represented by hyperechoic lines. B: unaffected limb, forearm point. C: affected limb, arm point; fat infiltration is visible, surrounded by hyperechoic lines in a circular arrangement. D: unaffected limb, arm point

0.5 to 0.8 moderate, and above 0.8 large (Cohen, 2009). Results were presented with their respective confidence intervals and considered statistically significant at $P < .05$ for both the analysis of findings and their association with temperatures.

3. RESULTS

A total of 121 women were screened, of whom 43 met the inclusion criteria as women who had unilateral breast cancer and completed all their treatment. In the laboratory, a thermogram was obtained for each participant. From these thermograms, four regions of interest (ROIs) were analyzed at the arm and forearm points—two in the affected upper limb and two in the unaffected upper limb—totaling 172 ROIs. Subsequently, four ultrasound images were generated for each participant, corresponding to the same points—two in the affected upper limb

and two in the unaffected upper limb—resulting in 172 images (Fig. 3).

The women included in the sample had a mean age of 54.14 years. Less than half of the sample (44.18%) presented lymphedema according to the volumetric evaluation. Among the clinical symptoms that may indicate the onset or presence of lymphedema, the most prevalent were a sensation of heaviness (67.45%), paresthesia (55.82%), and peau d'orange (44.19%) (Table 1).

The qualitative findings observed in the ultrasound images of the upper limb with and without lymphedema (Table 2) indicated that, at the forearm point (TA), fibrosis was highly prevalent (47.4%), with individuals with lymphedema being 9.9 times more likely to present this dermal alteration ($P = .01$). Thickening of the dermo epidermal layer was

observed in 57.9% of cases, with individuals with lymphedema having a 5.3 times higher likelihood of presenting this alteration ($P = .02$).

At the arm point (TB), fibrosis (42.1%) and dermo epidermal layer thickening (68.5%) were the most prevalent alterations. Among these, thickening of the dermo epidermal layer had the highest odds ratio, with individuals with lymphedema being 23.9 times more likely to present this alteration ($P < .001$). This was the sole dermal alteration with a statistically

significant association in the evaluated region of interest.

The comparison of the mean maximum, mean, and minimum temperatures obtained through thermography with the qualitative findings identified via ultrasound (Table 3) showed that, in general, the mean temperatures recorded for all findings at the corresponding points on the arm and forearm ranged from 26.46°C to 30.41°C in the limb affected by lymphedema, with moderate standard deviations. The effect size,




Fig. 3. Flowchart of sample screening and acquisition of ultrasound images and thermograms

Table 1. Sociodemographic and clinical characteristics of the sample (n = 43)

Variable	Mean (SD) or n (%)
Age in years (mean, SD)	54.14 (7.72)
Presence of lymphedema n (%)	19 (44.18)
Treatments n (%)	
Axillary lymphadenectomy	40 (93.02)
Adjuvant radiotherapy in axillary area	41 (95.34)
Symptoms n (%)	
Feeling of heaviness	29 (67.45)
Paresthesia (numbness)	24 (55.82)
Increased local temperature	14 (32.56)
Erythema (redness)	8 (18.61)
Peau d'orange (skin thickening resembling orange peel)	19 (44.19)
Presence of Lymphedema (>200ml) n (%)	
Without Lymphedema	24 (55.81)
With lymphedema	19 (44.19)

SD: standard deviation; n: sample size

Table 2. Tissue alterations in upper limbs identified in ultrasound images

Variable	With lymphedema (n=19) n (%)	Without lymphedema (n=24) n (%)	P value	Risk ratio (95% CI)
TA				
Fibrosis	9 (47.4)	2 (8.4)	.01	9.9 (1.8–54.5)
Fat infiltration	5 (26.3)	0	.05	18.6 (0.9–361.3)
DEC thickening	11 (57.9)	5 (20.8)	.02	5.3 (1.4–20.0)
Loss of differentiation	6 (31.6)	1 (4.17)	.04	10.7 (1.2–98.1)
TB				
Fibrosis	8 (42.1)	4 (16.7)	.07	3.7 (0.9–14.9)
Fat infiltration	4 (21.1)	0	.08	14.3 (0.8–283.0)
DEC thickening	13 (68.5)	2 (8.4)	<.001	23.9 (4.2–136.0)
Loss of differentiation	3 (15.8)	0	.13	10.4 (0.6–214.8)

P values were calculated using the Chi-square test; DEC: dermoepidermal layer; TA: forearm assessment point; TB: arm assessment point

calculated using Hedges' *g*, was small for the following conditions: fat infiltration at the minimum temperature ($g = 0.31$) at TA; mean temperature ($g = 0.46$) and minimum temperature ($g = 0.44$) at TB. Similarly, for loss of differentiation at TA, the mean temperature ($g = 0.23$) and minimum temperature ($g = 0.23$) showed small effect sizes.

None of the analyzed variables presented statistically significant differences between groups ($P > .05$), suggesting the absence of significant temperature variations associated with the qualitative ultrasound findings.

4. DISCUSSION

The ultrasound findings for the upper limb with lymphedema revealed a higher prevalence of dermoepidermal layer thickening (57.9%) and fibrosis (47.4%). The lymphedema group demonstrated the highest risk for fat infiltration in the forearm (18.6) and dermoepidermal thickening in the arm (23.9). However, these results were not statistically significant ($P > .05$) for fat infiltration at TA and TB, with only dermoepidermal thickening at TB showing significance.

Early identification of lymphedema is crucial to minimize its impact on quality of life and daily

functioning. Studies suggest that initial symptoms tend to decrease and stabilize 18 months post-surgery, while volume and circumference changes continue to increase up to 36 months post-surgery—key indicators of established lymphedema (Armer, 2019). In the present study, 29 women reported a sensation of heaviness in the limb, exceeding the number of women diagnosed with lymphedema (19). This finding suggests the potential for an early stage of lymphedema in the symptomatic sample, as heaviness and paresthesia are predictive indicators for early lymphedema intervention, as outlined in post-breast cancer treatment surveillance programs (Wong, 2024).

Some studies have reported that volumetry, a commonly used clinical method to assess lymphedema, is less effective because it cannot differentiate between tissue changes and fluid accumulation. This distinction is critical for determining the stage of lymphedema (Park et al., 2024; Rezende et al., 2023). Imaging methods, particularly ultrasound, have therefore gained prominence due to their ability to distinguish between tissue layers and identify the most affected areas of the upper limb. Literature shows that the anterior forearm (TA) is the region most frequently presenting tissue alterations, consistent with the findings of this study (Suehiro, 2016).

Table 3. Correlation between tissue alterations found in ultrasound and maximum, mean, and minimum temperatures of upper limbs

Tissue Alteration	TA			TB		
	Tmax	Tmean	Tmin	Tmax	Tmean	Tmin
	Mean (SD)	Mean (SD)	Mean (SD)	Mean (SD)	Mean (SD)	Mean (SD)
Fibrosis						
Present	28.92 (2.52)	29.69 (0.90)	27.83 (1.36)	30.23 (2.10)	29.85 (1.40)	28.78 (1.92)
Absent	28.05 (3.97)	29.88 (1.61)	27.73 (2.47)	29.49 (2.56)	30.34 (1.42)	28.98 (2.08)
<i>P</i> value	.48	.57	.67	.44	.41	.70
Hedges' <i>g</i>	-0.13	0.10	0.08	-0.14	0.16	0.07
Fat Infiltration						
Present	29.48 (2.54)	29.34 (1.71)	26.46 (2.81)	29.62 (2.24)	29.35 (1.21)	27.82 (1.07)
Absent	28.08 (3.87)	29.89 (1.52)	27.83 (2.31)	29.59 (2.53)	30.32 (1.42)	29.01 (2.07)
<i>P</i> value	.39	.49	.24	.92	.12	.13
Hedges' <i>g</i>	-0.23	0.19	0.31	0.03	0.46	0.44
Dermo epidermal Layer Thickness						
Present	28.69 (2.67)	29.34 (1.59)	27.21 (2.41)	30.15 (2.28)	29.94 (1.68)	28.49 (1.86)
Absent	28.04 (4.04)	29.98 (1.50)	27.87 (2.33)	29.48 (2.54)	30.35 (1.36)	28.49 (2.08)
<i>P</i> value	.62	.23	.28	.40	.58	.27
Hedges' <i>g</i>	-0.08	0.19	0.17	-0.14	0.09	0.18
Loss of Differentiation						
Present	29.12 (2.43)	29.28 (1.24)	26.82 (2.63)	31.53 (2.97)	30.76 (1.19)	28.66 (2.16)
Absent	28.08 (3.91)	29.91 (1.55)	27.83 (2.32)	29.52 (2.47)	30.26 (1.43)	28.97 (2.06)
<i>P</i> value	.50	.31	.31	.15	.49	.83
Hedges' <i>g</i>	-0.15	0.23	0.23	-0.49	-0.24	0.07

TA: forearm assessment point; TB: arm assessment point; Tmax: maximum temperature; Tmin: minimum temperature; Tmean: mean temperature; SD: standard deviation

Another significant result of this study is the high prevalence of dermoepidermal layer thickening in both regions of interest. This aligns with evidence that, across all stages of lymphedema involving volume changes, dermal edema emerges as the first and most characteristic alteration associated with the pathology (Ricci et al., 2022).

Ultrasound evaluation in this study also identified a greater number of women with fibrosis in the affected limb. Literature suggests that subcutaneous and skin changes in lymphedema are caused by extracellular alterations, such as connective tissue hypertrophy (Carvalho et al., 2020). Fibrosis is more common from stage 2 of lymphedema, while fat accumulation typically arises in stage 3 (Bowman & Rockson, 2024). This indicates that most participants in the current study had moderate lymphedema, as only a small number of individuals exhibited fat infiltration, while a larger proportion presented with fibrosis and dermoepidermal thickening.

Thermography has been increasingly utilized for the diagnosis of lymphedema, showing promise as a non-invasive and complementary tool. Recent studies have demonstrated its reproducibility and accuracy in aiding

lymphedema diagnosis, particularly in breast cancer survivors (Debiec-Bąk et al., 2020; Gomes, 2024a, 2024b). Thermography has been shown to detect initial temperature elevations in the limb that reflect ongoing inflammatory processes, with evidence supporting its good-to-excellent reproducibility across various postures and regions of interest. However, in the present study, no significant association was observed between thermographic findings and tissue alterations. This suggests that thermography may lack the sensitivity required to differentiate between clinical stages of lymphedema and their associated tissue changes. Further studies with larger and more diverse samples are needed to validate these findings and to explore thermography's potential role in clinical staging and monitoring of lymphedema.

However, this study concluded that thermography has limitations in sensitivity for detecting tissue alterations, as no significant correlation was found with ultrasound findings. It is important to note that the sample was heterogeneous, with regions of interest often showing multiple tissue alterations, complicating the precise correlation of temperature with individual findings. The lack of statistical

significance in the correlated data is primarily attributed to the small sample size, despite the presence of effect size (Hedges' *g*), which was small in some analyses.

5. CONCLUSION

The study identifies a significant prevalence of tissue alterations in individuals with lymphedema, particularly fibrosis and thickening of the dermoepidermal layer, which were notably more common in affected upper limbs. Participants with lymphedema demonstrated significantly higher ratio of these alterations, with statistically significant associations observed at both arm and forearm assessment points. Conversely, thermographic analysis did not reveal meaningful temperature differences that correlated with the ultrasound findings, indicating its limited effectiveness in detecting these specific alterations. These findings reinforce the value of ultrasound imaging as a reliable tool for assessing dermal changes in upper limbs affected by lymphedema, while suggesting that thermography requires further refinement to enhance its diagnostic utility.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

CONSENT

All authors declare that 'written informed consent was obtained from all participants.

ETHICAL APPROVAL

All authors hereby declare that all experiments have been examined and approved by the appropriate ethics committee under protocol no. 5.434.586 and have therefore been performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki.

ACKNOWLEDGEMENT

The authors thank the Universidade Federal de Pernambuco and CAPES for the financial support provided for the translation of the manuscript and payment of publication fees. The authors also thank FACEPE for financial support

for this study (APQ 0801-4.08/21 and APQ-1330-4.08/21).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Armer, J. M., et al. (2019). Lymphedema symptoms and limb measurement changes in breast cancer survivors treated with neoadjuvant chemotherapy and axillary dissection: Results of American College of Surgeons Oncology Group (ACOSOG) Z1071 (Alliance) substudy. *Support Care Cancer*, 27, 495–503.
- Bowman, C., & Rockson, S. G. (2024). The role of inflammation in lymphedema: A narrative review of pathogenesis and opportunities for therapeutic intervention. *International Journal of Molecular Sciences*, 25(7), 3907.
- Carvalho, V. L. de, Pitta, G. B. B., & Cunha, S. X. S. (2020). Uso de software de imagem ultrassonográfica para diferenciação de edema de origem venosa e de origem linfática em membros inferiores. *Jornal Vascular Brasileiro*, 19, e20190139.
- Cohen, J. (2009). *Statistical power analysis for the behavioral sciences* (2nd ed.). Psychology Press.
- Dębiec-Bąk, A., et al. (2020). Using thermography in the diagnostics of lymphedema: Pilot study. *Lymphatic Research and Biology*, 18(3), 247–253.
- Denlinger, C. S., et al. (2018). Survivorship, version 2.2018, NCCN clinical practice guidelines in oncology. *Journal of the National Comprehensive Cancer Network*, 16(10), 1216–1247.
- Donahue, P. M. C., et al. (2023). Advances in the prevention and treatment of breast cancer-related lymphedema. *Breast Cancer Research and Treatment*, 200(1), 1–14.
- Estupiñán, A. I., et al. (2020). Correlation between indocyanine green lymphography and thermography to evaluate areas of dermal backflow in lymphedema. *Journal of Plastic, Reconstructive & Aesthetic Surgery*, 73(10), 1897–1916.
- Executive Committee, et al. (2016). The diagnosis and treatment of peripheral lymphedema: Consensus document of the

- International Society of Lymphology. *Lymphology*, 49(4), 170–184.
- Fernández-Cuevas, I., et al. (2017). Termografia infravermelha para a detecção de lesões em medicina esportiva. In Quesada, J. P. (Ed.), *Application of Infrared Thermography in Sports Science* (pp. 217–231). Springer International Publishing.
- Gomes, V. M. d. S. A., et al. (2024a). Accuracy of infrared thermography in diagnosing breast cancer-related lymphedema. *Journal of Clinical Medicine*, 13(20), 6054.
- Gomes, V. M. d. S. A., et al. (2024b). Reproducibility of thermography for measuring skin temperature of upper limbs in breast cancer survivors. *Biomedicines*, 12(11), 2465.
- Goudarzi, S., et al. (2023). Segmentation of arm ultrasound images in breast cancer-related lymphedema: A database and deep learning algorithm. *IEEE Transactions on Biomedical Engineering*, 70(9), 2552–2563.
- Kelly-Hope, L. A., et al. (2021). Infrared thermal imaging as a novel non-invasive point-of-care tool to assess filarial lymphoedema. *Journal of Clinical Medicine*, 10(11), 2301.
- Keszytüs, D., et al. (2023). Use of infrared thermography in medical diagnosis, screening, and disease monitoring: A scoping review. *Medicina*, 59(12), 2139.
- Levenhagen, K., et al. (2017). Diagnosis of upper quadrant lymphedema secondary to cancer: Clinical practice guideline from the oncology section of the American Physical Therapy Association. *Physical Therapy*, 97(7), 729–745.
- Mander, A., et al. (2019). Upper limb secondary lymphedema ultrasound mapping and characterization. *International Angiology: A Journal of the International Union of Angiology*, 38(4), 334–342.
- Park, J. Y., Jeon, J. Y., & Cha, S. (2024). Ultrasonography. *Ultrasonography*, 43(4), 284–293.
- Polat, A. V., et al. (2020). Efficacy of ultrasound and shear wave elastography for the diagnosis of breast cancer-related lymphedema. *Journal of Ultrasound in Medicine*, 39(4), 795–803.
- Rezende, L. F. de, et al. (2023). Ultrassonografia como instrumento de avaliação do linfedema secundário ao câncer de mama: Revisão sistemática. *Jornal Vascular Brasileiro*, 22, e20220144.
- Ricci, V., et al. (2022). From physical to ultrasound examination in lymphedema: A novel dynamic approach. *Journal of Ultrasound*, 25(1), 1–7.
- Shavit, E., Wollina, U., & Alavi, A. (2018). Lipoedema is not lymphoedema: A review of current literature. *International Wound Journal*, 15(6), 921–928.
- Shen, A., et al. (2023). Risk factors of breast cancer-related lymphoedema: Protocol of an umbrella review. *BMJ Open*, 13(4), e070907.
- Suehiro, K., et al. (2016). Skin and subcutaneous tissue ultrasonography features in breast cancer-related lymphedema. *Annals of Vascular Diseases*, 9(4), 312–316.
- Wong, H. C. Y., et al. (2024). Multinational Association of Supportive Care in Cancer (MASCC) clinical practice guidance for the prevention of breast cancer-related arm lymphoedema (BCRAL): International Delphi consensus-based recommendations. *EClinicalMedicine*, 68, 101393.
- Yusof, K., et al. (2021). Assessment of potential risk factors and skin ultrasound presentation associated with breast cancer-related lymphedema in long-term breast cancer survivors. *Diagnostics*, 11(8), 1303.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/128074>