



Annual Research & Review in Biology

Volume 39, Issue 12, Page 199-206, 2024; Article no.ARRB.128844

ISSN: 2347-565X, NLM ID: 101632869

(Past name: Annual Review & Research in Biology, Past ISSN: 2231-4776)

The Story of the Pink Bollworm, *Pectinophora gossypiella* (Saunders) on Cotton

Rakesh S^{a*}, Udikeri S S^a, Mahamed Ashiq I^b and Akarsh S V^c

^a Department of Agricultural Entomology, College of Agriculture, University of Agricultural Sciences, Dharwad-580005, Karnataka, India.

^b Department of Biotechnology, College of Agriculture, University of Agricultural Sciences, Dharwad-580005, Karnataka, India.

^c Department of Agronomy, Keladi Shivappa Nayaka University of Agricultural and Horticultural Sciences, Shivamogga-577412, Karnataka, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/arrb/2024/v39i122182>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/128844>

Review Article

Received: 24/10/2024

Accepted: 28/12/2024

Published: 30/12/2024

ABSTRACT

Cotton, a vital commercial fibre crop, is prone to heavy infestations by insect pests, with bollworms being the most damaging globally. *Helicoverpa armigera* and other Heliothines have developed significant resistance to conventional insecticides, rendering cotton cultivation uneconomical in many regions by the mid-1990s. The introduction of Bt cotton, expressing Cry1Ac toxins, marked a breakthrough in bollworm management, drastically reducing pesticide usage and increasing yields. However, resistance to Cry toxins, particularly Cry1Ac, emerged over time, as first reported in pink bollworm (*Pectinophora gossypiella*) in Gujarat, India, by 2010. Factors contributing to resistance

*Corresponding author: E-mail: mragriinfo@gmail.com;

include mutations in genes such as *PgCad1*, ABC transporters, and others. Additionally, declining toxin expression in Bt cotton hybrids and improper pest management practices exacerbate resistance. Integrated pest management strategies, such as timely sowing, field sanitation, pheromone traps, and mating disruption tools, have been recommended to mitigate resistance. Recent data show an increase PBW infestations in North India, which is being driven by factors like shorter cotton seasons and little prior exposure to Bt toxins. Studies focusing on genetic and phenotypic polymorphism in Indian PBW populations have emphasized the need for sustainable resistance management. Adopting dual-toxin Bt cotton (Cry1Ac + Cry2Ab) and stringent IRM strategies remains critical to preserving the efficacy of transgenic technology.

Keywords: Cotton; *bacillus thuringiensis*; cry toxin; pink bollworm; resistance.

1. INTRODUCTION

Cotton is an important commercial fibre crop which is attacked by many types of insect pests throughout the season. Among these, the complex of American, pink, spotted, and spiny bollworms are said to be the most dreaded leading to heavy loss and requiring huge protection costs worldwide (Rajendran, et al., 2018, Garg, et al., 2022). Cotton is attacked in the United States of America by *Heliothis virescens* (Fabricius) and *Helicoverpa zea* (Boddie) (Blanco, 2012), in India by *Helicoverpa armigera* (Hübner), and in Australia by *H. armigera* and *H. punctigera* Wallengren (Downes & Mahon, 2012). Out of the cost of cotton production in India, a lion part of more than 43% was speared targeting insect pest management, of which 80%for bollworm control itself, particularly *H. armigera* (Kranthi, 2012). Heliothines caused a significant threat to upland cotton production worldwide including India in the late 1980s due to their high degree of resistance to classical insecticides (organophosphates, carbamates and pyrethroids) (Razaq, 2019). In India, *H. armigera* showed 15-fold resistant to quinalphos, 5 to 6500-fold to cypermethrin, >30 fold to methomyl, 16 to 3200-fold to fenvalerate, and 2 to 59-fold to endosulfan. Similarly, *Earias vittella* Fabricius also exhibited 70 to 362 folds resistance to these insecticides (Armes, et al., 1996). Hence by the mid-1990s cotton cultivation mostly turned uneconomical in major cotton growing countries (Kranthi, 2012, Narayanamoorthy & Kalamkar, 2006) particularly in the Indian and China. The use of insecticides having novel chemistry was too costly to rely upon. Meanwhile, efforts to popularize Integrated Pest Management (IPM) strategies to manage bollworms remained enforcement activity rather than self-adaption by farmers (Kranthi, 2012). Hence as the best-bet technique to manage bollworms, MON 531 (Cry1Ac gene) event-based *Gossypium hirsutum* varieties DP20B, DP50B, DP90B, NuCotn 33, and NuCotn 35 were planted

successfully in the USA on a commercial scale during 1996 (Perlak, et al., 2001). By 2000, *Bt* cotton genotypes were approved in subsequently, other countries like Australia (1996), Argentina (1997), China (1997), Mexico (1998), South Africa (1998), Indonesia (2001) and Colombia (2002) (Steven, et al., 2008). In all of these countries *H. armigera*, *H. zea*, *H. virescens*, *Earias* spp, and pink bollworm, *Pectinophora gossypiella* (Saunders) were successfully controlled. This enforced India also to accept transgenic technology by 2002 (Mohan & Manjunath, 2002) and three *G. hirsutum* Bt cotton hybrids (MECH 12, MECH 162, and MECH 184) were authorized for commercial cultivation (Khadi, 2007). Due to great bioefficacy against bollworms, the *Bt* cotton area increased drastically from 0.29 lakh ha in 2002-03 to 33.53 lakh ha in 2006-07 (Anonymous, 2020). In this period success of *Bt* transgenic was characterized by cent per cent bollworm check, a 40-60%reduction in pesticide usage in cotton (Khadi, 2007), and net profit enhanced to 78% (Rao and Dev, 2009).

The resistance development to cry1Ac is an expected phenomenon from many of the laboratory studies show no change in the susceptibility during the first 4-5 generations of *H. armigera* under selected pressure against Cry1Ac but, initial indications of resistance were clear after the 6th selection regimen and by the end of 10th generation, resistance increased 76-folds as reflected by the LC₅₀ values (Kranthi, et al., 2000). Hence, to avoid/slowdown resistance development to Cry1Ac toxin in filed population growing of refugia imposed by EPA, US and all other Government agencies including GEAC in India, as resistance management strategies. Another IRM technique for sustainability of transgenic technology under the threat potential of resistance development was keeping refugia also in use (Andow, 2008). This was the cultivation of genotypes that expressed two toxins i.e., Cry1Ac and Cry2Ab simultaneously.

Cry1Ac + Cry2Ab (MON 15985) cotton was progressively replacing Cry1Ac varieties (DP20B, DP50B, DP90B, NuCotn 33 and NuCotn 35) in the United States during 2003 and NuCotn 37, Deltapine varieties in Australia during 2004. The replacement was spurred by the idea that evolution of resistance would be delayed substantially by two-toxins relative to one toxin (Ali & Luttrell, 2007). In India also during 2006, Bollgard II was launched which has two genes, Cry1Ac + Cry2Ab with unrelated mode of action that aids in delaying resistance. By 2013, Bollgard II occupied 91% of *Bt* cotton (Choudhary & Gaur, 2010) and by 2015, more than 95% of *Bt* cotton with stacked genes (Kranthi, 2016, Komarlingam, 2018). However, the major gene in action is cry1Ac itself in all of these genotypes (Anonymous, 2003). So far, over more than two decades field resistance to cry toxin has not been observed in any Heliothines including *H. armigera* in any region of the world. But in 2010, pink bollworm, *P. gossypiella* showed resistance to Bollgard I event (Cry1Ac) in Gujarat, with a resistance ratio of 44 (Dhurua & Gujar, 2011) and subsequently to Bollgard II (Cry1Ac + Cry2Ab) with 40-80% of the bolls harboured surviving larvae as recorded from Amreli and Bhavnagar districts of Gujarat during 2014 (Naik, et al., 2018). From 2014 onwards, reports have highlighted the outbreak of pink bollworm on Bollgard I and Bollgard II in states namely, Gujarat, Madhya Pradesh, Maharashtra, Karnataka, and Andhra Pradesh with a resistance ratio of 1387 to Cry1Ac and 4196 to Cry2Ab in 2017 from central and southern India (Naik et al., 2018). The PBW is essentially a monophagous pest, relying exclusively on cotton as its primary host, with no significant alternate hosts in Indian cotton ecosystems (Ingram, 1994). This fact was continued through a systematic survey in Karnataka (Rakesh, et al., 2023). PBW primarily feeds on developing seeds. The rapid and widespread adoption of *Bt* cotton, produced substantial quantities of *Cry* toxins in raw seeds (Kranthi, et al., 2005) and bolls (Knight, et al., 2013), leading to high selection pressure causing resistance. The exchange of seeds and movement of seed cotton allow resistant populations to spread from one location to another (Naik, et al., 2020).

The resistance development might be due to various reasons

- Cry toxin expression is highest in leaves followed by squares, bolls, and flowers (Kranthi, et al., 2005, Willrich, et al., 2009, Mahalakshmi & Prasad, 2013).

- Toxin expression decline after 90 days from sowing (Iqbal, et al., 2013, Bhullar & Gill, 2015).
- Feeding of segregating seeds in F₁ *Bt*-cotton hybrids (Shahid, et al., 2021).
- Non-application of refugia (Shahid, et al., 2021).
- Extending crop life (Fand, et al., 2019, Krishna et al., 2020).
- Lack of timely and appropriate management initiatives (Krishna, et al., 2020).
- Hybrids have different flowering and fruiting periods (Fand, et al., 2019, Krishna, et al., 2020).
- Cultivation of long duration hybrids (Fand, et al., 2019).
- Long term storage of raw cotton in ginneries (Kumar, et al., 2020).
- Nitrogen deficits and condition reduced the levels of toxin expression (Kranthi, 2012).

This resulted in more damage (40-95 %) with 20-30% yield loss, despite the use of novel chemical insecticides and integrated pest management strategies by the farmers (Fand, et al., 2019).

Following field scale survival many efforts have been made to know the resistance level in Indian PBW populations. The flower infested PBW from Junagadh, Amreli, Jalna, Wardha, Yavatmal, and Rajkot showed 3.6, 26.4, 3.2, 5.6, 1.4, 3.4- and 3.6-fold resistance to Cry1Ac, respectively; Adilabad, Prakasham, Wardha, Yavatmal and Jalna population showed 28.67, 18.67, 67.67, 20.67 and 630.67-fold resistance to Cry2Ab, respectively. Whereas, the boll infested PBW from Rajkot (Gujarat) recorded 371.8-fold to Cry1Ac and 4214.3-fold resistance to Cry2Ab (Naik, et al., 2021). Similarly, Cry toxin resistance levels in pink bollworm across cotton-growing zones in India is represented in Table 1. Previous studies reported no incidence of pink bollworm (PBW) on *Bt* cotton in North India (Naik, et al., 2018; Shantala, 2020). However, in recent years, the severity of green boll infestation by PBW has been increasing in the northern cotton-growing zones. The infestation levels have ranged from 10 to 100 percent in Punjab, with an average larval incidence of 25.20 %, and in Haryana, with 27.90 % larval incidence. In Rajasthan, infestation levels ranged from 10 to 30 percent, with an average larval incidence of 11.20 % (Prasad & Kumar, 2022, Sham, 2023, Rakesh, 2024). The lower adaptability of PBW to Cry toxins in Northern India could be attributed

Table 1. Cry toxin resistance levels in pink bollworm across cotton-growing zones in India

Cry toxins	Cotton growing zones			Year	Authors
	Southern	Central	Northern		
Resistance ratio against Cry1Ac (-folds)	18-47	18-121	Nil	2013	Naik, et al., 2018
	39	36-767	Nil	2014	
	36-551	13-1581	Nil	2015	
	19-343	36-2868	Nil	2016	
	798	704-2060	Nil	2017	
	-	372	-	2020	Naik, et al., 2021
	711-817	419-740	719-1042	2021	Sham, 2023
	386-858	535-960	306-856	2022	Rakhesh, 2024
	1	2-31	Nil	2013	Naik, et al., 2018
	1	3-423	Nil	2014	
Resistance ratio against Cry2Ab (-folds)	15-1604	36-2465	Nil	2015	Naik, et al., 2021
	192-5003	287-3779	Nil	2016	
	1570	1306-9366	Nil	2017	
	-	4214	-	2020	
	613-784	335-907	871-1038	2021	
	809-1584	970-2096	630-1225	2022	

to several factors. Notably, PBW populations in the region had not been exposed to BG-I cotton prior to the simultaneous introduction of both BG-I and BG-II for commercial cultivation in 2006. Additionally, the cotton-growing season in North India is relatively short, lasting only five to six months to accommodate subsequent wheat cultivation. This creates a closed season, significantly reducing selection pressure by limiting the number of PBW generations exposed to Bt cotton each year (Naik, et al., 2018).

Mutations in certain genes are the primary cause of Cry toxin resistance in the pink bollworm (*Pectinophora gossypiella*). The *PgCad1* gene, which encodes cadherin proteins, often exhibits mutations or deletions that reduce or eliminate the binding of Cry1Ac toxin to midgut receptors, a critical step for the toxin's action. Similarly, mutations in ATP-binding cassette (ABC) transporter genes impair the transport or activation of Cry toxins, which contributes to resistance. Alterations in aminopeptidase N (APN) genes further disrupt the binding of Cry toxins, compounding the resistance mechanism. Additionally, mutations in the alkaline phosphatase (ALP) gene, which is associated with receptor function, can hinder the effectiveness of Cry toxins in the insect gut. These genetic changes collectively enable the pink bollworm to withstand the toxic effects of Cry proteins (Tabashnik, et al., 2004, Morin, et al., 2004, Malthankar & Gujar, 2015, Agrawal, et al., 2020, Shanthala, 2020, Fabrick, et al., 2021). In addition to conventional diet incorporation bioassays used to assess resistance levels,

efforts have been made to investigate Cry toxin resistance polymorphism in Indian pink bollworm (PBW) strains. However, these studies have predominantly focused on sequencing the mitochondrial *COI* gene (Naik, et al., 2020). This has served as evidence for resistant population expansion. The Cry toxin resistance in PBW has been linked to certain mutations in cadherin genes and this is being validated presently in respect of Indian strains (Shanthala, 2020, Sham, 2023, Rakhesh, 2024).

Integrated pest management practices (Rakhesh, et al., 2023):

- Timely sowing and narrowing sowing window (Henneberry, et al., 1982).
- Implement field sanitation, clearing old stalks, and eliminating unopened and partly opened bolls (Gutierrez, et al., 2015).
- Select verified *Bt* cotton seeds for planting.
- Monitoring of PBW using trap catches (8-10 moths for 3 consecutive days, Qureshi et al., 1993) and 10% flower/ green boll damage (ETL).
- Use pheromone traps to mass trapping of pink bollworm (>20 traps/ha) at 45 DAS.
- Spraying of Ovicides at 60 DAS (Udikeri, et al., 2022).
- Two release of *T. bactre* @ 60000 per acre between 75-85 DAS (Udikeri, et al., 2022).
- Use of mating disruption tools (SPLAT/ PB rope L).
- Synthetic pyrethroid spray after 100 DAS (Gopalaswamy, et al., 2000).

- Timely crop termination (Khakwani, et al., 2022)

2. CONCLUSION

Cotton is a crucial commercial crop, but it faces constant threats from a wide range of insect pests, with the bollworms viz., American, pink, spotted and spiny bollworms being among the most destructive. These pests cause significant yield losses and incur high control costs globally. In India, the primary bollworm species affecting cotton crops are *Helicoverpa armigera* and *Pectinophora gossypiella*, with resistance to chemical pesticides becoming a major concern since the late 1980s. This resistance, especially to organophosphates, carbamates, and pyrethroids, led to economic losses in cotton production, particularly in major growing countries like India and China. The introduction of Bt cotton, with its Cry1Ac toxin, revolutionized bollworm management by offering a more sustainable control method. Bt cotton significantly reduced pesticide use and increased profits, leading to its widespread adoption. However, the emergence of resistance in pink bollworm populations, particularly in Gujarat, raised concerns. Resistance to Cry1Ac and Cry2Ab toxins has been observed in various cotton-growing regions, highlighting the need for robust resistance management strategies. Efforts to mitigate resistance include the development of Bt cotton varieties expressing multiple toxins, such as Cry1Ac and Cry2Ab, to delay the evolution of resistance. Integrated pest management (IPM) practices, such as timely sowing, field sanitation, the use of pheromone traps, and crop termination, are also essential for maintaining the efficacy of Bt cotton. Despite these challenges, ongoing research and management strategies remain critical for the sustainability of cotton production in India.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Agrawal, A., Venkatesan, T., Ramasamy, G. G., Ramesan Syamala, R., Muthugounder, M., & Rai, A., (2020). Transcriptome alterations of field-evolved resistance in *Pectinophora gossypiella* against Bt Bollgard II cotton in India. *Journal of Applied Entomology*, 144(10), 929-940.
- Ali, M. I., & Luttrell, R. G. (2007). Susceptibility of bollworm and tobacco budworm (*Lepidoptera: Noctuidae*) to Cry2Ab2 insecticidal protein. *Journal of Economic Entomology*, 100(3), 921-931.
- Andow, D. A. (2008). The risk of resistance evolution in insects to transgenic insecticidal crops. *Collection of Biosafety Reviews*, 4, 142-199.
- Anonymous, (2003). *Bacillus thuringiensis* Cry2Ab2 protein and the Genetic Material Necessary for its Production in Cotton (006487) Factsheet. (https://www.epa.gov/opp00001/biopesticides/ingredients/factsheets/factsheet_006487.htm)
- Anonymous, (2020). Area, production and productivity of GM Crops, Ministry of Agriculture & Farmers Welfare, PIB, Delhi. www.pib.gov.in/PressReleasePage.aspx?PRID=1605056
- Armes, N. J., Jadhav, D. R., & De Souza, K. R. (1996). A survey of insecticide resistance in *Helicoverpa armigera* in the Indian subcontinent. *Bull. Entomol. Res.*, 86, 499-514.
- Bhullar, H. S., & Gill, R. S. (2015). Cry toxin expression on different plant parts and crop stages among transgenic Bt cotton hybrids. *Pest. Res. J.*, 27, 155-159.
- Blanco, C. A. (2012) *Heliothis virescens* and Bt cotton in the United States. *GM crops and food*, 3(3), 201-212.
- Choudhary, B., & Gaur, K. (2010). Bt cotton in India: A country profile. *ISAAA Series of Biotech Crop Profiles*. ISAAA: Ithaca, New York, pp. 1-32.
- Dhurua, S., & Gujar, G. T. (2011). Field-evolved resistance to Bt toxin Cry1Ac in the pink bollworm, *Pectinophora gossypiella* (Saunders) (*Lepidoptera: Gelechiidae*), from India. *Pest Management Science*, 67, 898-203.
- Downes, S., & Mahon, R. (2012). Evolution, ecology and management of resistance in *Helicoverpa* spp. to Bt cotton in Australia. *Journal of invertebrate pathology*, 110(3), 281-286.

- Fabrick, J. A., LeRoy, D. M., Mathew, L. G., Wu, Y., Unnithan, G. C., Yelich, A. J., Carrière, Y., Li, X., & Tabashnik, B. E. (2021). CRISPR-mediated mutations in the ABC transporter gene ABCA2 confer pink bollworm resistance to Bt toxin Cry2Ab. *Scientific reports*, 11(1), 10377.
- Fand, B. B., Nagrare, V. S., Gawande, S. P., Nagrale, D. T., Naikwadi, B. V., Deshmukh, V., Narkhedkar, G. N., & Waghmare, V. N. (2019). Widespread infestation of pink bollworm, *Pectinophora gossypiella* (Saunders) (Lepidoptera: Gelechiidae) on Bt cotton in Central India: a new threat and concerns for cotton production. *Phytoparasitica*, 47(3), 313-325.
- Garg, R., Singh, B., Kargwal, R., Tiwari, S., Yadav, S., Jakhar, A., Kumar, D., & Sharma, B. L., (2022) Resistance in Pink Bollworm *Pectinophora gossypiella* (Saunders) against Bt Cotton, a Major Threat to Cotton in India: A Brief Review. *International Journal of Plant and Soil Science*, 34(22): 248-61.
- Gutierrez A P, Ponti L, Herren H R, Baumgärtner J and Kenmore P E, 2015, Deconstructing Indian cotton: Weather, yields, and suicides. *Environmental Sciences Europe*, 27(1): 1-17.
- Gopalswamy S V S, Rao N H P and Hanumanth Rao V, 2000, Insecticides in the control of pink bollworm in cotton. *Pestology*, 24(7): 7-11.
- Henneberry T J, Kittock D L and Bariola L A, 1982, Pink bollworm: effect of cotton types and planting date on early season infestations. *Southwestern Entomologist*, 7: 9-65.
- Ingram, W. R. (1994). *Pectinophora* (Lepidoptera: Gelechiidae) in Insect pests of cotton. Wallingford: CAB International, pp. 107-149.
- Iqbal, A., Ali, S., Zia, M. A., Shahzad, A., Din, J. U., Asad, M. A. U., Ali, G. M., & Zafar, Y. (2013). Comparative account of Bt gene expression in cotton under normal and salt affected soil. *International Journal of Agriculture and Biology*, 15(6), 1181-1186.
- Khadi, B. M. (2007). Success Story of Bt Cotton in India. In an *International Cotton Advisory Council Meeting at Burkina Faso, December*.
- Khakwani K, Cengiz R, Naseer S, Asif M and Sarwar G, 2022, Cotton pink bollworm (*Pectinophora gossypiella*) management with the goal of eradication from the cotton producing countries of the world. *Applied Ecology and Environmental Research*, 20(2): 1199-1213.
- Knight, K., Head, G., & Rogers, J. (2013). Season-long expression of Cry 1Ac and Cry 2Ab proteins in Bollgard II cotton in Australia. *Crop Protection*, 44, 50-58.
- Komarlingam, M. S. (2018). Refuge-in-bag for Bt cotton. *Current Science*, 114, 726-731.
- Kranthi, K. R. (2012). Bt cotton-questions and answers. *Indian Society for Cotton Improvement (ISCI), Mumbai*.
- Kranthi, K. R. (2016). Possible implications of a recent gazette notification on Bt cotton scenario in India. *Current Science*, 111, 1588-1590.
- Kranthi, K. R., Kranthi, S., Ali, S., & Banerjee, S. K. (2000). Resistance to CryIAc δ -endotoxin of *Bacillus thuringiensis* in a laboratory selected strain of *Helicoverpa armigera* (Hubner). *Current Science*, 78(8), 1001-1004.
- Kranthi, K. R., Naidu, S., Dhawad, C. S., Tatwawadi, A., Mate, K., Patil, E., Bharose, A. A., Behere, G. T., Wadaskar, R. M., & Kranthi, S. (2005). Temporal and intra-plant variability of Cry1Ac expression in Bt-cotton and its influence on the survival of the cotton bollworm, *Helicoverpa armigera* (Hübner) (Noctuidae: Lepidoptera). *Current Science*, 89(2), 291-298.
- Krishna, M. S., Reddy, Y. R. & Chandrayudu, E. (2020). Validation of pink bollworm *Pectinophora gossypiella* (Saunders) management strategies in Bt cotton. *Journal of Entomology and Zoology Studies*, 8(5), 2064-2067.
- Kumar, R., Monga, D., Naik, V., Singh, P., & Waghmare, V. N. (2020) Incipient infestations and threat of pink bollworm *Pectinophora gossypiella* (Saunders) on Bollgard-II cotton in the northern cotton-growing zone of India. *Current Science*, 118(9), 1454-1456.
- Mahalakshmi, M. S., & Prasad, N. V. (2013). Seasonal expression of Cry 1 Ac toxin in different plant parts of two Bt Cotton hybrids. *Annals of Plant Protection Sciences*, 21(1), 62-64.
- Malthankar, P.A., & Gujar, G.T. (2015). Association of Cry2Ab resistance with alkaline phosphatase in pink bollworm *Pectinophora gossypiella* (Saunders) (Lepidoptera: Gelechiidae). *Biopestic. Int*, 10, 143-151.

- Mohan, K. S., & Manjunath, T. M., (2002). *Bt* cotton-India's first transgenic crop. *Journal of Plant Biology-New Delhi*, 29(3), 225-236.
- Morin, S., Henderson, S., Fabrick, J. A., Carrière, Y., Dennehy, T. J., Brown, J. K., & Tabashnik, B. E. (2004). DNA-based detection of *Bt* resistance alleles in pink bollworm. *Insect biochemistry and molecular biology*, 34(11), 1225-1233.
- Naik, V. C. B., Subbireddy, K. B., Kranthi, S., Nagrare, V. S., Kumbhare, S., Narkhedkar, G. N., & Waghmare, V. N. (2021). Pink bollworm, *Pectinophora gossypiella* (Saunders) (Lepidoptera: Gelechiidae) survival on transgenic cotton in India. *Egyptian Journal of Biological Pest Control*, 31(1), 1-7.
- Naik, V. C., Kumbhare, S., Kranthi, S., Satija, U., & Kranthi, K. R. (2018). Field-evolved resistance of pink bollworm, *Pectinophora gossypiella* (Saunders) (Lepidoptera: Gelechiidae), to transgenic *Bacillus thuringiensis* cotton expressing crystal 1Ac (Cry1Ac) and Cry2Ab in India. *Pest Management Science*, 74(11), 2544-2554.
- Naik, V., Pusadkar, P. P., Waghmare, S. T., Raghavendra, K. P., Kranthi, S., Kumbhare, S., Nagrare, V. S., Kumar, R., Prabhulinga, T., Narkhedkar, G. N., & Waghmare, V. N. (2020). Evidence for population expansion of cotton pink bollworm *Pectinophora gossypiella* (Saunders) (Lepidoptera: Gelechiidae) in India. *Scientific reports*, 10(1): 1-11.
- Narayanamoorthy, A., & Kalamkar, S. S. (2006). Is *Bt* cotton cultivation is economically viable for Indian farmers-An empirical analysis. *Economic and Political Weekly*, 41(26), 2716-2724.
- Perlak, F. J., Oppenhuizen, M., Gustafson, K., Voth, R., Sivasupramaniam, S., Heering, D., Carey, B., Ihrig, R. A. & Roberts, J. K. (2001). Development and commercial use of Bollgard® cotton in the USA-early promises versus today's reality. *The Plant Journal*, 27(6), 489-501.
- Prasad, Y. G., & Kumar, R. (2022). Pink bollworm in North cotton growing zone of India-It is time for coordinated corrective measures. Mumbai, India: Weekly Publication of Cotton Association of India, *Cotton Statistics and NEWS*. 3, 1-8.
- Rajendran, T. P., Birah, A., & Burange, P. S. (2018). Insect pests of cotton. In *Pests and their management*. Springer, 361-411.
- Rakshesh, S. (2024). Evaluation of diverse strains of pink bollworm, *Pectinophora gossypiella* (Saunders) for cry toxin resistance and its validation through markers. *Ph.D. (Agri.) Thesis, University of Agricultural Sciences, Dharwad, Karnataka, India*.
- Rakshesh, S., Anil, K. M., Krishna N. L., & Santhosha, K. M., (2023) A brief review on integrated pest management of pink bollworm, *Pectinophora gossypiella* (Saunders) in cotton. *Insect Environment*, 26(3):375-387. DOI: 10.55278/YWCK6622
- Rakshesh, S., Hanchinal, S. G., Bheemanna, M., Hosamani, A. K., & Nidagundi, J. M. (2023). Incidence of pink bollworm, *Pectinophora gossypiella* (Saunders) on various malvaceous plants in North Eastern region of Karnataka, India. *Journal of Farm Sciences*, 36(3), 266-269. DOI:10.61475/JFS.2023.v36i3.06
- Rao, N. C. & Dev, S. M. (2009). Socio-economic impact of transgenic cotton. *Agricultural Economics Research Review*, 22(347-2016-16863), 461-470.
- Razaq, M., Mensah, R., & Athar, H. U. R. (2019). Insect pest management in cotton. *Cotton production*, pp. 85-107.
- Shahid, M. R., Farooq, M., Shakeel, M., Ashraf, M., Zia, Z. U., Ahmad, S., & Mahmood, A. (2021). Need for growing non-*Bt* cotton refugia to overcome *Bt* resistance problem in targeted larvae of the cotton bollworms, *Helicoverpa armigera* and *Pectinophora gossypiella*. *Egyptian Journal of Biological Pest Control*, 31(1), 1-8.
- Sham, S. G. (2023). Studies on the genetics of resistance in Cry toxin resistant populations of pink bollworm, *Pectinophora gossypiella* (Saunders) (Lepidoptera: Gelechiidae) and its management. *Ph.D Thesis, University of Agricultural Sciences, Raichur, Karnataka, (India)*.
- Shanthala, P. K. (2020). Analysis of biochemical and molecular basis of cry toxin resistance in pink bollworm, *Pectinophora gossypiella* (Saunders) infesting *Bt* cotton. *M. Sc. (Agri.) Thesis, University of Agricultural Sciences, Dharwad, Karnataka, India*.
- Steven, N. E., Ruberson, J. R., Sharma, H. C., Wilson, L., & Wu, K. (2008). The present and future role of insect-resistant genetically modified cotton in IPM. In *Integration of insect-resistant genetically modified crops within IPM programs*, Springer, pp. 159-194.
- Tabashnik, B. E., Liu, Y. B., Unnithan, D. C., Carrière, Y., Dennehy, T. J., & Morin, S.

- (2004). Shared genetic basis of resistance to *Bt* toxin Cry1Ac in independent strains of pink bollworm. *Journal of Economic Entomology*, 97(3), 721-726.
- Udikeri S S, Siddalingappa Hugar, Kuberappa Gundannavar, Poornima Holeyannavar and Rajesab Nadaf, 2022, Development and validation of an adaptable IPM module for pink bollworm in BG-II *Bt* transgenic cotton. Book of abstracts World Cotton Research Conference 7, Cairo, Egypt, 3-7, October 2022. pp. 477-485.
- Willrich, S. M., Patterson, T. G., Gilles, G. J., Nolting, S. P., Braxton, L. B., Leonard, B. R., Van, D. J. W., & Lassiter, R. B. (2009). Quantification of Cry1Ac and Cry1F *Bacillus thuringiensis* insecticidal proteins in selected transgenic cotton plant tissue types. *Journal of Economic Entomology*, 102(3), 1301-1308.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/128844>