



Regeneration Status of Different Conifers in Gulmarg Forest Range of Kashmir Himalayas

**Suraya Tasveer ^a, Aasif Ali Gatoo ^{a*}, Akhlaq Amin Wani ^a,
M.A. Islam ^a, Shah Murtaza Mushtaq ^a and P. A. Khan ^b**

^a Division of NRM, Faculty of Forestry, Benhama, Ganderbal, SKUAST-Kashmir-191201, India.

^b Division of FBT, Faculty of Forestry, Benhama, Ganderbal, SKUAST-Kashmir-191201, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/arja/2025/v18i1641>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/129426>

Original Research Article

Received: 05/11/2024

Accepted: 07/01/2025

Published: 11/01/2025

ABSTRACT

The present study was concentrated in the four forest blocks, namely Khaipora, Tangmarg, Ferozpora and Baderkut of Gulmarg Forest Range in Kashmir Himalayas. Natural regeneration and edaphic factors affecting regeneration status of different conifers was studied. Quadrats of 2x2m in the main plot of size 50x20m were laid down to study the regeneration at four blocks of Gulmarg Forest Range. The maximum numbers of recruits (current year seedlings, less than height of 8 cm), unestablished (height more than 8cm to 200cm) and established (height more than 200cm) plants were found for *Abies pindrow* at all the sites with the exception at Ferozpora where unestablished plants of *Pinus wallichiana* were found highest. Highest establishment stocking per cent and regeneration success was found for *Abies pindrow* at Khaipora, Ferozpora and Baderkut. Overall regeneration success was found to be highest at Baderkut followed by Khaipora while as the lowest regeneration success was found at Ferozpora respectively. Simple correlation study reveal that the

*Corresponding author: E-mail: asifskuast@gmail.com;

regeneration success has a significant positive relationship with Light Intensity, pH, Electric conductivity, organic carbon, available nitrogen and available potassium while as litter layer and available phosphorous had a significant negative correlation with regeneration success.

Keywords: Natural regeneration; regeneration success; edaphic factors; conifers.

1. INTRODUCTION

The Himalayan Forest ecosystem plays a substantial role in providing many ecological services to the human population and their livestock; however due to inadequate and unreliable information, management of these forest types have not been addressed suitably. Most watersheds in the Himalayan region are experiencing a decline in different forest covers and agricultural land use type has happened to be the main constituent of the current landscape (Sundriyal *et al.*, 1994). In India, conifers are one of the most valuable natural resources which contribute significantly to its socio-economic development by providing goods and services to the people and industries. They play a major role in enhancing the quality of environment by influencing the basic life support system and generate considerable revenue. In Western Himalayas, the Himalayan moist temperate forest, extending from 1500-3000 m amsl is of immense significance from the environmental conservation and sustainable development (Sharma and Baduni, 2000). Although qualitative descriptions of the forest vegetation of the Himalayas are available (Champion and Seth, 1968), scattered attempts on quantitative examinations have also been made (Saxena and Singh, 1982; 1984; Ali *et al.*, 2009). Regeneration is a key process for the survival of species in a community under diverse environmental conditions (Walkley & Black, 1954). Assessment of regeneration status of plant communities has a paramount importance for sustainable conservation and management. Natural regeneration is essential as it addresses mainstream biodiversity concerns. Therefore, it is necessary to pay attention to the quantitative and qualitative factors and environmental conditions to study the regeneration status in a forest (Karami *et al.*, 2017).

Inadequate regeneration is the main problem of forests in mountain regions (Krauchii *et al.*, 2000). Sustainable conservation of forests involves proper planning and management of seedlings, saplings and young trees that ensure maintenance of forest community structure and ecological stability (Moravie *et al.*, 1997). Successful management of forest covers need reliable research data on aspects such as structural attributes and demographic profile of tree species. Therefore, it is important to study the regeneration potentials of different natural forest trees in Kashmir Himalaya to determine the probable trends of vegetation in the future.

2. MATERIALS AND METHODS

Study area: The study area is in Gulmarg forest range of Jammu and Kashmir Himalayas. Gulmarg forest range is divided into seven blocks out of which we conducted study in four blocks viz; Khaipora, Ferozpora, Tangmarg and Baderkut in during 2019-2020. The Gulmarg Forest range is situated between 34°3' 31.2" N to 74° 23' 1.0" E. The total area of Gulmarg forest range is 15933.78 ha. The procedure of Stratified random Sampling was followed.

Regeneration assessment: The regeneration status of different conifers was recorded on the basis of the number of individuals occurring at seedling, sapling and pole stage in each quadrat. The height of unestablished, established plants and recruits were measured for the assessment of regeneration (Champion, 1935). The data collected was analysed using the following formulae which was given by Chacko (1965).

Table 1. Details of the study area

Block	Block Name	Elevation (m amsl)	Longitude	Latitude	Aspect	Slope
1	Khaipora	2345	34°03'15.9"N	74°24'40.0"E	NW	35°
2	Tangmarg	2265	34°03'40.93"N	74°25'15.58"E	NE	35°
3	Ferozpora	2365	34°2'55.59"N	74°28'50.98"E	SW	45°
4	Baderkut	2221	34°02'25.64"N	74°27'04.29"E	NE	45°

$$\begin{aligned}\text{Recruits (r)/ha} &= 2500 \sum_{i=1}^n r_i / m \\ \text{Unestablished regeneration (u)/ha} &= 2500 \sum_{i=1}^n u_i / m \\ \text{Established regeneration (e) /ha} &= 2500 \sum_{i=1}^n e_i / m\end{aligned}$$

Where

- n - Number of sampling units
- m – Total number of recording units in survey
- ri – Total number of recruits in each sampling unit
- ui – Total number of unestablished plants in each sampling unit
- ei – Total number of established plants in each sampling unit

$$\text{Weighted Average height (m)} = \frac{\text{Total height of unestablished regeneration} + (\text{Number. of established plants} \times \text{establishment height})}{\text{Total unestablished plants} + \text{total established plants}}$$

On the basis of above estimates following indices will be calculated:

$$\begin{aligned}\text{Establishment index (I}_1\text{)} &= \frac{\text{Weighted average height}}{\text{Establishment height}} \\ \text{Stocking index (I}_2\text{)} &= 1/2500 \times \frac{\text{Unestablished regeneration/ha}}{4} + \text{Established regeneration/ ha}\end{aligned}$$

$$\text{Established stocking per cent} = 100 (I_1 \times I_2)$$

$$\text{Regeneration success (\%)} = \text{Stocking index (I}_2\text{)} \times 100$$

Edaphic factors: Following parameters were studied for accessing the edaphic factors affecting natural regeneration of different conifers

- Organic matter layer
- Physico-chemical properties

Organic matter layer (cm): It was measured as depth of the column from top of humus layer to the point under humus where soil exists.

Physico-chemical properties: Mridaparikshak – A mini lab developed by ICAR-Indian institute of soil science (IISS), Bhopal in 2015 will be used for estimation of soil organic carbon (Walkley and Black, 1954), Soil PH (Jackson, 1973), EC, available Nitrogen (Subbiah and Asija, 1956), available Phosphorus and available Potassium (Jackson, 1973).

3. RESULTS AND DISCUSSION

The present investigation for regeneration studies were carried out on the basis of recruits, unestablished, established and regeneration

success at four different blocks namely Khaipora, Tangmarg, Ferozpora Baderkut in Gulmarg Forest Range. The data of various regeneration compounds are presented in Table 2 which are described below:

The regeneration ability of any species is a vital factor in determining its ability to produce its offsprings and is ruled by number of factors (Seildling & Constien, 1998). The natural regeneration of a tree species largely depends on the production and germination of seeds and the establishment of seedlings and saplings Rao (1998). The pattern of population dynamics of seedlings, saplings and adults of a species can reveal the recruitment profile, which is used to define their regeneration status. The results recorded pertaining to natural regeneration are interpreted and correlated to the literature to understand the regeneration status of conifers in different blocks of Gulmarg forest range. The present study revealed a total of four conifer species. Regeneration status was analyzed using the methodology of Chacko (1965).

The perusal of data in Table 2 and Table 3 reveal that natural regeneration of conifers were present at all the blocks except for *Taxus contorta* which was present only in Khaipora and *Cedrus deodara* which was present at Tangmarg and Baderkut. The presence of recruits, unestablished and established plants of all the conifers were present at their respective sites with the exception of *Cedrus deodara* which were absent in Tangmarg but present in Baderkut. But in case of *Taxus contorta* established plants could not be found. Of all the species found in Khaipora among recruits, unestablished and established, percentage of *Abies pindrow* was higher (55.74%). Similarly, percentage of *Pinus wallichiana* was higher (46.15%) in Tangmarg. Among the Species that were found in Ferozpora percentage of *Abies pindrow* was higher (40.00%). In Baderkut percentage of *Picea smithiana* was higher than other species (50.00%). The regeneration of species varied at different elevations which are in accordance with the study conducted by Gupta (1996). The lesser number of recruits, un-established and established regeneration at lower altitude may be attributed to high biotic interference (grazing, trampling by the animals and human settlement). The lesser number of recruits, un-established and established regeneration at lower altitude may be attributed to high biotic interference (grazing, trampling by the animals and human settlement). Similar trend was reported by Mir (2016) while working on *Betula* forest in Northwest Himalayas, Lankar et al. (2010) & Lankar, (2007), while working on Natural regeneration status of the endangered medicinal plant, *Taxus baccata*, in northwest Himalaya. Similarly, Table 3 reveals that the maximum weighted average height (271.71), Establishment index (1.35), Establishment stocking per cent (9.06) was obtained at Ferozpora, while as Maximum Stocking Index (0.26) was obtained at Baderkut. The results obtained are within the range of Gatoo (2021) who obtained similar range while working on Natural regeneration status of *Betula* in Sangla valley of Indian Himalayas, Gupta (2019) while working on Natural regeneration status of Indian hazelnut in western Himalaya of Himachal Pradesh. Lower values of weighted average height, Establishment index, Stocking Index, Establishment stocking per cent might be attributed to the overexploitation of the species for timber, it could also be attributed to lower incidence of solar influx in the area, as supported

by the findings of Niemann (1992) and Seidling and Constein (1998) who investigated the importance of light for yew survival. Highest regeneration success (26.58 %) was recorded at Baderkut and the minimum regeneration success (19.81) was recorded at Ferozpora (Table 3). Among species the highest regeneration success was obtained for *Abies pindrow* at Khaipora (11.22 %) and at Ferozpora (7.27). whereas maximum regeneration success was obtained for *Pinus wallichiana* (9.78) at Tangmarg block and for *Picea smithiana* (14.29 %) at Baderkut. The results were in conformity with the findings of Sharma (2006), Gupta (2007), Lanker (2007) and Malik et al. (2012). The regeneration of trees generally depends upon the ability of trees to provide sufficient quantity of seeds, their ability to germinate and grow as seedlings and survive in the under-canopy environment, where soil moisture and light may often be limiting Kozlowski (1971), Good and Good (1972). A number of factors measured to be responsible for absence or poor natural regeneration such as thick layer of humus/depth of organic matter layer Gordon (1970), Troup (1921), Parnell (1930), Taylor et al. (1934), Glover (1936) Mahendru (1936) Dhillon (196) Kaushik (1954), Datta (1958), Kaul (1970) Sufi (1970) Rattan (2011) continuous grazing Redcliffe (1906) Troup (1921) Flewett (1930) Yadav (1963) Sufi (1970) and low germination capacity of the seeds.

It is evident from Table 4 that regeneration success has a significant positive correlation with Light intensity (0.75), pH (0.71), Electric conductivity (0.85), Organic carbon (0.94), available nitrogen (0.99) and available potassium (0.76) while as Organic matter layer (-0.97) and available phosphorous (-0.81) has a significant negative correlation. The findings from our study got support from the findings of Colaona and Giannini (1971), Singh (1983) and Pengshalin et al. (2006), where they have reported that depth of organic matter has indirect relationship with regeneration success. The findings also got support from Puri (1955) while studying the Himalayan conifer. Simple correlation coefficient results are in conformity with the finding of Filipiak and Komisaraek (2005) who reported potassium content and organic carbon in soil has positive effect on the regeneration of silver fir and Gatoo et al. (2020) who reported organic carbon, pH, solar influx, nitrogen and potassium in soil has a positive effect of regeneration of Silver fir and *Betula utilis*.

Table 2. Existing regeneration status of different tree species in different sites of Gulmarg Forest Range

Species	Recruits/ha		Unestablished/ ha		Established/ ha		Density/ha
	Number	Per cent (%)	Number	Per cent (%)	Number	Per cent (%)	
Khaipora							
<i>Picea smithiana</i> (Wall.)Boiss.	1250.00±37.50	31.39	373.75±13.17	28.85	74.25±3.44	33.11	200.00
<i>Abies pindrow</i> Spach	1986.67±80.83	49.88	622.92±21.95	48.08	125.00±3.75	55.74	280.00
<i>Pinus wallichiana</i> A.B.Jacks.	498.33±17.56	12.51	249.17±8.78	19.23	25.00±0.75	11.15	100.00
<i>Taxus contorta</i> Griff.	247.50±11.46	6.21	49.83±1.76	3.85	0.00±0.00	0.00	60.00
Total	3982.50	100.00	1295.67	100.00	224.25	100.00	640.00
Tangmarg							
<i>Abies pindrow</i> Spach	1000.00±30	37.38	250.83±6.29	32.26	100.33±2.52	30.77	200.00
<i>Pinus wallichiana</i> A.B.Jacks.	1300.00±39	48.60	376.25±9.44	48.39	150.50±3.77	46.15	160.00
<i>Picea smithiana</i> (Wall.)Boiss.	375.00±11.25	14.02	150.50±3.77	19.35	75.25±1.89	23.08	180.00
<i>Aesculus Indica</i> (Wall.ex Cambess.)	0.00± 0.00	0.00	0.00±0.00	0.00	0.00±0.00	0.00	80.00
<i>Robinia pseudoacacia</i> L.	0.00±0.00	0.00	0.00±0.00	0.00	0.00±0.00	0.00	60.00
<i>Cedrus deodara</i> (Roxb.ex D.Don) G.Don	0.00±0.00	0.00	0.00±0.00	0.00	0.00±0.00	0.00	100.00
Total	2675.00	100.00	777.58	100.00	326.08	100.00	780.00
Ferozpora							
<i>Pinus wallichiana</i> A.B.Jacks.	551.83±13.84	19.47	200.67±5.03	42.11	125.42±3.15	33.33	160.00
<i>Abies pindrow</i> Spach	2006.67±50.33	70.80	125.42±3.15	26.32	150.50±3.77	40.00	200.00
<i>Picea smithiana</i> (Wall.)Boiss.	275.92±6.92	9.73	150.50±3.77	31.58	100.33±2.52	26.67	180.00
Total	2834.42	100.00	476.59	100.00	376.25	100.00	540.00
Baderkut							
<i>Pinus wallichiana</i> A.B.Jacks.	275.92±6.92	11.58	100.33±2.52	9.52	50.17±1.26	12.50	100.00
<i>Picea smithiana</i> (Wall.)Boiss	602.00±15.10	25.26	326.08±8.18	30.95	150.50±3.77	37.50	160.00
<i>Abies pindrow</i> Spach	1505.00±37.75	63.16	627.08±15.73	59.52	200.67±5.03	50.00	180.00
<i>Cedrus deodara</i> (Roxb.ex D.Don) G.Don	0.00±0.00	0.00	0.00±0.00	0.00	0.00±0.00	0.00	100.00
Total	2382.92	100.00	1053.49	100.00	401.34	100.00	440.00

Table 3. Establishment and stocking data for different tree species in different sites at gulmarg forest range

Species	Weighted average height (cm)	Esatblishment Index(I1)	Stocking Index (I2 final)	Established Stocking Per cent (I1xI2x100)	Regeneration Success
Khaipora					
<i>Picea smithiana</i> (Wall.)Boiss.	36.36	0.18	0.06	1.21	6.70
<i>Abies pindrow</i> Spach	36.87	0.18	0.11	2.07	11.22
<i>Pinus wallichiana</i> A.B.Jacks.	21.62	0.10	0.03	0.37	3.49
<i>Taxus contorta</i> Griff.	2.50	0.01	0.0005	0.006	0.49
Total	97.36	0.48	0.21	3.67	21.92
Tangmarg					
<i>Abies pindrow</i> Spach	59.91	0.29	0.06	1.95	6.52
<i>Pinus wallichiana</i> A.B.Jacks.	59.90	0.29	0.09	2.92	9.78
<i>Picea smithiana</i> (Wall.)Boiss.	69.58	0.34	0.04	1.57	4.51
<i>Aesculus Indica</i> (Wall.ex Cambess.)	0.00	0.00	0.00	0.00	0.00
<i>Robinia pseudoacacia</i> L.	0.00	0.00	0.00	0.00	0.00
<i>Cedrus deodara</i> (Roxb.ex D.Don) G.Don	0.00	0.00	0.00	0.00	0.00
Total	189.39	0.94	0.20	6.45	20.81
Ferozpora					
<i>Pinus wallichiana</i> A.B.Jacks.	78.96	0.39	0.07	2.77	7.02
<i>Abies pindrow</i> Spach	110.68	0.55	0.07	4.02	7.27
<i>Picea smithiana</i> (Wall.)Boiss.	82.07	0.41	0.05	2.26	5.51
Total	271.71	1.35	0.19	9.06	19.81
Baderkut					
<i>Pinus wallichiana</i> A.B.Jacks.	68.59	0.34	0.03	1.03	3.01
<i>Picea smithiana</i> (Wall.)Boiss.	66.38	0.33	0.09	3.08	9.28
<i>Cedrus deodara</i> (Roxb.ex D.Don) G.Don	0.00	0.00	0.00	0.00	0.00
<i>Abies pindrow</i> Spach	51.40	0.25	0.14	3.67	14.29
Total	186.34	0.93	0.26	7.78	26.58

Table 4. Site characteristics of different conifer forest stands at Gulmarg Forest Range

Site name	Regeneration success	Light Influx (%)	Litter layer (cm)	pH	Electric Conductivity	Organic Carbon (%)	Nitrogen (kg/ha)	Phosphorus (kg/ha)	Potassium (kg/ha)
Khaipora	21.92	28.08	1.70	6.8 ^a	0.90	0.35%	218.70	22.34	13.00
Ferozpura	19.81	19.36	4.50	6.4	0.70	0.25%	175.00	36.64	10.00
Baderkut	26.58	28.78	0.50	6.8 ^a	0.94	0.41%	287.50	20.00	22.35
Tangmarg	20.81	16.79	3.80	6.5	0.75	0.29%	200.00	28.60	19.34
CD	0.34	0.36	0.45	0.091	0.0135	0.0094	3.32	0.40	0.36

4. CONCLUSION

The maximum numbers of recruits, unestablished and established plants were found for *Abies pindrow* at all the sites with the exception at however unestablished plants of *Pinus wallichiana* were found to be highest at Ferozpora. Highest establishment stocking per cent and regeneration success was found for *Abies pindrow* at Khaipora, Ferozpora and Baderkut while as for *Pinus wallichiana* at Tangmarg. Overall regeneration success was found to be highest at Baderkut followed by Khaipora while as the lowest regeneration success was found at Ferozpora respectively. The main factors responsible for poor regeneration seem to be higher organic matter layer and available phosphorus.

Simple correlation study reveal that the regeneration success has a significant positive relationship with Light Intensity, pH, Electric conductivity, organic carbon, available nitrogen and available potassium while as litter layer and available phosphorous has a significant negative correlation with regeneration success.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

ACKNOWLEDGEMENT

Authors could never accomplish this task without the help of so many generous people. Authors would like to acknowledge Dean, Faculty of Forestry SKUAST-K J&K, HOD and other staff members of Division of NRM FOF SKUAST-K J&K for their help, support, advices, instruction, assistance and patience.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

Ali, A., Shamet, G. S., & Kumar, R. (2009). Evaluation of natural regeneration status and site quality characteristics of *Taxus wallichiana* forest in Himachal Pradesh. *Annals of Biology*, 25(2), 159-162.

- Chacko, V. J. (1965). *Manual of sampling technique for forest surveys*. Manager Publications.
- Champion, H. G., & Seth, S. K. (1968). *A revised survey of forest types of India*. Manager Publications.
- Champion, H. G. (1935). *Silvicultural research manual for use in India*. Vol. I. *The experimental manual*. Government of India Press.
- Colaona, M., & Giannini, R. (1971). Seed germination and early growth of *Abies alba*, *Epilobium angustifolium*, and *Rubus hirtus* as influenced by various natural seed beds. *Annals Accademia Italians di Science Forestall*, 20, 227-258.
- Datta, R. C. (1958). Natural regeneration of spruce and fir. In *Proceedings of First Regional Meeting of Silviculturists, Himalayan Region*, Kashmir, F. R. I. & Dehradun, 13-15.
- Dhillon, G. S. (1961). Recent trends in natural regeneration technique of silver fir and spruce forests. In *Proceedings of Tenth Silvicultural Conference*, 1, 313-321.
- Flewett, W. E. (1930). The reproduction of spruce and fir forests. In *Proceedings of Punjab Forestry Conference*, 64-66.
- Filipiak, M., & Komisarek, J. (2005). Regeneration of the European silver fir (*Abies alba* Mill.) in the Sudety Mountains on soils with different physico-chemical properties. *Dendrobiology*, 53, 17-25.
- Gatoo, A. A., Wani, A. A., Islam, M. A., Murtaza, S., Bhat, A., & Zehra, S. (2021). Natural regeneration status of *Betula utilis* in Sangla valley of Indian Himalayas. *Journal of Pharmacognosy and Phytochemistry*, 9(2), 1574-1580.
- Glover, H. M. (1936). The practical problem of the management of the Himalayan fir forests. *Indian Forester*, 62(5), 276-282.
- Good, N. F., & Good, R. E. (1972). Population dynamics of tree seedlings and saplings in mature eastern hardwood forests. *Bulletin of the Torrey Botanical Club*, 99, 172-178.
- Gordon, D. T. (1970). Natural regeneration of white and red fir. U.S. Forest Research Paper, Paaf, 5th West Forest Range Experiment Station, PSW 58, 32.
- Gupta, D., Tiwari, C., Rajput, B. S., Kumar, S., & Sharma, D. (2019). Natural regeneration status of Indian hazelnut (*Corylus jacquemontii* - Decne.) in Western Himalaya of Himachal Pradesh. *Journal of Pharmacognosy and Phytochemistry*, 8(1), 327-331.

- Gupta, D. (2007). Regeneration status and growth distribution in silver fir and spruce forest. M.Sc. Thesis, Dr. Y. S. Parmar University of Horticulture and Forestry, Solan, India.
- Gupta, N. K. (1996). Appraisal of vegetational pattern of Shimla district through remote sensing with special reference to the ecology of fir-spruce forests. Ph.D. Thesis, Dr. Y. S. Parmar University of Horticulture and Forestry, Solan, India.
- Jackson, M. L. (1973). *Soil chemical analysis*. Prentice Hall of India Pvt. Ltd.
- Karami, A., Karamshahi, A., & Shahi, E. (2017). Effects of forestry practices on the regeneration and biodiversity of woody plants in the northern forest ecosystems of Iran. *Geology, Ecology and Landscapes*, 1(4), 21-36.
- Kaul, P. N. (1970). Natural regeneration of silver fir and spruce. In *Proceedings of State Forest Conference, J&K State*, Srinagar: Forest Department, J & K State, 79-84.
- Kaushik, R. C. (1954). Management of spruce (*Picea morinda*) and fir (*Abies pindrow*) and *Abies webbiana* forests. *Indian Forester*, 80(2), 79-89.
- Kozlowski, T. T. (1971). *Growth and development of trees. Vol. I: Seed germination, ontogeny and shoot growth*. Academic Press.
- Krauchi, N., Brang, P., & Schonenberger, W. (2000). Forests of mountainous regions: Gaps in knowledge and research needs. *Forest Ecology and Management*, 132, 73-82.
- Lankar, U., Malik, A. R., Gupta, N. K., & Butola, J. S. (2010). Natural regeneration status of the endangered medicinal plant, *Taxus baccata* Hook. f. syn. *T. wallichiana* in North-West Himalaya. *International Journal of Biodiversity Science, Ecosystem Services and Management*, 6(1-2), 20-27.
- Lankar, U. (2007). Studies on some edapho-ecological characteristics and regeneration status of Himalayan yew (*Taxus baccata* Hook. f. syn. *T. wallichiana* Zucc.).
- Mahendru, I. D. (1936). The problem of natural regeneration of silver fir (*Abies spectabilis*). *Indian Forester*, 62(5), 261-275.
- Malik, A. R., Shamet, G. S., & Butola, J. S. (2012). Natural regeneration status of Chilgoza pine (*Pinus gerardiana* Wall.) in Himachal Pradesh, India: An endangered pine of high edible value. *Applied Ecology and Environmental Research*, 10(3), 365-373.
- Mir, N. A. (2016). *Community structure, regeneration, and soil nutrient status of Betula utilis D. Don stands in high altitude forests of central and North Kashmir* (Ph.D. thesis). Sheri-Kashmir University of Agricultural Science and Technology, Jammu and Kashmir, India.
- Moravie, M. A., Pascal, J. P., & Auger, P. (1997). Investigating canopy regeneration process through individual-based spatial models: Applications to tropical rain forest. *Ecological Modeling*, 104, 241-260.
- Niemann, H. (1992). Preservation of natural occurrences of yew. *AFZ. Allgemeine Forst Zeitschrift*, 47(8), 405-407.
- Parnell, R. (1930). The reproduction of spruce and fir forests. In *Proceedings of Punjab Forestry Conference*, Lahore, 57-58.
- Pengshalin, Wang-Chang, W., Liming, G., Wenjain, L., Faliang, & Chen, W. (2006). Major ecological factors affecting growth of *Taxus* spp. *Acta Scientiarum Naturalium Universitatis Sunyatsani*, 45(3), 65-69.
- Puri, G. S., & Mohan, N. P. (1955). The Himalayan conifers III. The succession of forest communities oak-conifers of the Bushar Himalaya. *Indian Forester*, 81, 405-487.
- Rao, G. R. (1998). Study on dynamics of herbage layer in pine and khair-based natural silvipastoral system in Northwest Himalaya. Ph.D. Thesis, Dr. Y. S. Parmar UHF, Solan, H.P.
- Redcliffe, E. (1906). Research on the regeneration of silver fir. *Indian Forester*, 32(8), 402-404.
- Saxena, A. K., & Singh, J. S. (1982). A phytosociological analysis of woody species in forest communities of a part of Kumaun Himalaya vegetation. *Plant Ecology*, 50, 3-22.
- Seildling, W., & Constien, A. (1998). Spontaneous yew regeneration in Grunewald. *AFZ Der Wald, Allgemeine Forst Zeitschrift für Waldwirtschaft und Umweltvorsorge*, 53(21), 1318-1320.
- Sharma, C. M., & Baduni, N. P. (2000). Effect of aspect on the structure of some natural stands of *Abies pindrow* in Himalayan moist temperate forest. *Springer*, 20, 309-317.
- Sharma, Y. (2006). *Studies on stand parameters and natural regeneration status of silver fir and spruce in Himachal Pradesh* (M.Sc. thesis). Dr. Y.S. Parmar UHF, Solan, H.P.
- Singh, R., Thakur, G. C., & Sood, V. P. (1983). Phytosociology and resource utilization by

- different forest trees in southeastern slopes around Shimla, Himachal Pradesh. *Indian Forester*, 120(2), 1108-1119.
- Subbiah, B. V., & Asija, G. L. (1956). A rapid procedure for the estimation of available nitrogen in soils. *Current Science*, 25, 259-260.
- Sufi, G. R. (1970). Regeneration of silver fir and spruce in western Himalayas with specific reference to the state of Jammu and Kashmir. In *Proceedings of State Conference, J&K*, 85-93.
- Sundriyal, R. C., Sharma, E., Rai, L. K., & Rai, S. C. (1994). Tree structure, regeneration, and woody biomass removal in a subtropical forest of Mamlaym watershed in Sikkim Himalaya. *Vegetation*, 113, 53-63.
- Taylor, E. M., Mehta, M. L., & Hoon, R. C. (1934). A study of the soils in the hill areas of Kulu Forest Division, Punjab, Part I. An investigation of soil profile under *Deodar*, spruce, blue pine, and chir. *Indian Forest Records (New Series) Silviculture*, 1(2), 289-346.
- Troup, R. S. (1921). *The silviculture of Indian trees* (Vol. III). Clarendon Press.
- Walkley, A. J., & Black, I. A. (1954). Estimation of soil organic carbon by chronic acid titration method. *Soil Science*, 37, 29-28.
- Yadav, J. S. P. (1963). Studies on soil profile in Chakrata division of Uttar Pradesh. *Indian Forester*, 89(1), 18-38.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2025): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/129426>