



Phytochemistry and Medicinal Uses of *Lonchocarpus* Species: Systematic Review of a Promising Medicinal Plant

**Sani M. Isyaka ^{a,b}, Olutola A. Olayemi ^c,
and Abdullahi M. Abdullahi ^{b,d*}**

^a Department of Chemical Sciences, North-Eastern University, Gombe, Nigeria.

^b Centre for African Medicinal Plants Research (CAMPRE), North-Eastern University, Gombe, Nigeria.

^c Department of Chemistry, Faculty of Science, University of Abuja, Nigeria.

^d Department of Chemistry, Federal University of Health Sciences Azare, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/sarjnp/2025/v8i1178>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/129910>

Review Article

Received: 11/11/2024

Accepted: 13/01/2025

Published: 15/01/2025

ABSTRACT

Research on natural products is crucial to the hunt for novel drugs to treat human diseases. Over time, there has been evidence and documentation of the usage of these medicinal plants as a source of physiologically active compounds with therapeutic potential for the treatment of a wide range of illnesses. Although the plant genus *Lonchocarpus* (*Fabaceae*, *papilionadeae*) has not yet been thoroughly studied, it has been shown to contain a wide range of compounds with numerous unique uses. This article discusses the pharmacological and biological roles, uses, and phytochemical constituents of the three species of *Lonchocarpus* plants such as; *Lonchocarpus*

*Corresponding author: Email: abdullahi.muhammadjika@fuhsa.edu.ng;

bussei, *Lonchocarpus cyanescens*, and *Lonchocarpus sericeus* that are found throughout Nigeria. A wide variety of polyphenolic compounds can be found in abundance in *Lonchocarpus*. Chalcones, flavans, aurones, dibenzoylmethane derivatives, rotenoids, pterocarpanes, flavanols, flavones, isoflavones, and flavonols are some of the classes that are included in this category, according to phytochemical studies. Despite the widespread need for medicinal compounds to treat both recognized and unknown ailments worldwide, little attention has been paid to the isolation, phytochemistry, and therapeutic qualities of compounds from *Lonchocarpus* species. Thus, the purpose of this review was to address the phytochemistry of *Lonchocarpus* species and to draw attention to the plant's therapeutic benefits and phytochemistry, particularly for those studying natural products and drug synthesis.

Keywords: *Flavonoids; isolation; Lonchocarpus; pharmacological; phytochemicals.*

1. INTRODUCTION

Because they are generally safe and have a positive effect on human health, specialists from all over the world have long been interested in plant-derived medications. In the pharmaceutical industry, plants having a long history of use in ethnomedicine can provide valuable medicines for the treatment of a wide range of illnesses and infectious diseases. It is believed that medicinal plants include most of the many kinds of bioactive compounds with a range of therapeutic properties. Over 80% of the world's developing population relies on traditional medicines as their main source of basic healthcare. According to the WHO, a range of medicinal plants can be used to make diverse medications (Chaachouay and Zidane, 2024; Fransworth et al., 1985).

Throughout human history, people have utilized natural remedies to alleviate illnesses (Abdullahi et al., 2007). This is primarily because conventional medicine has shown that these natural chemicals offer health advantages, making them attractive candidates for medication development (Cassidy et al., 2011). Consequently, novel medications continue to be made from plants (Cseke et al., 2006). A study by Moronkola et al., (2009) found that active ingredients or plant extracts from higher plants are present in 25% of the medications that pharmacists prescribe. Nowadays, it is believed that 119 chemical compounds originating from 90 distinct plant species are important drugs. 74% of the medications were found by examining the active components found in plants that were traditionally utilized for medical purposes.

Furthermore, 50% of the most often prescribed drugs in the US have a connection to natural goods or were motivated by them. Therefore, either the natural compounds themselves were employed directly or the phytochemicals were

used to create synthetic derivatives or analogs (Dorcas and Ibrahim, 2013). According to Vogler et al., (2006), 80% of people utilize medications derived from plants for medical purposes. Anticancer treatments and medications for infectious disorders are the most commonly utilized (Regnault-Roger and Calvo, 2004). There are still many higher plants that require investigation because only a small portion of those species have been examined for potential pharmacological application (Abdullahi et al., 2007).

Among these higher plants, the *Fabaceae* family's *Lonchocarpus* series is one whose thorough study is still lacking. According to Haegnauer and Grayer-Barkmaier (1993), the *Fabaceae* family, commonly known as *Leguminosae*, is the biggest family of flowering plants. One characteristic that sets plants in this family apart is the pod, or legume as it is known in science (Schrire et al., 2005). About 20,000 species are spread among 650 genera in this family, which is rich in terpenoids, polysaccharides, nitrogen-containing compounds, flavonoids, lipids, and anthraquinones (Wojciechowski et al., 2004; Xueqin et al., 2011). It is today one of the most commercially significant families in terms of the production of medicines, dye, fodder, decorations, wood, tannins, essential oils, resins, fragrances, piscicides, insecticides, and even human food (Kokwaro, 2009). Additionally, as legumes that fix nitrogen, they have an effect on the ecology (Bentjee, 1994; Mannetje, 2002).

Many of the fundamental medications used in conventional medicine are made from plants in the *Fabaceae* family. Alkaloids, terpenoids, flavonoids, steroids, phenylpropanoids, glycosides, phenols, and a variety of essential oils are among its well-known constituents, according to Xueqin et al. (2011). There are

about 100 species of *Lonchocarpus* (Fabaceae, Papilionoideae) in Madagascar, Africa, the Caribbean Islands, tropical America, and Australia (Magalhães et al., 2007; Mesía-Vela et al., 2001). According to chemical investigations, at least 35 of those species have been examined; these studies suggest that the plants contain compounds that may account for their usage in traditional medicine (Jain and Bari, 2010). Numerous *Lonchocarpus* species have been shown to be rich in polyphenolic compounds (Cassidy et al., 2011). Despite the widespread need for medicinal compounds to treat both recognized and unknown ailments worldwide, little attention has been paid to the isolation, phytochemistry, and therapeutic qualities of compounds from *Lonchocarpus* species. Thus, the purpose of this review was to address the phytochemistry of *Lonchocarpus* species and to draw attention to the plant's therapeutic benefits, particularly for those studying natural products and drug synthesis.

2. *Lonchocarpus*

The genus *Lonchocarpus* Fabaceae, Papilionoideae has over 100 species in tropical America, Africa, the Caribbean Islands, Australia, and Madagascar (Estrada-Castillón et al., 2024; Vogler et al., 2006; Mesía-Vela et al., 2001; Borges-Argáez et al., 2000; Jain and Bari, 2010; Lima et al., 2009). Chemical studies of at least 35 of those species suggest that they contain compounds that may account for their use in traditional medicine (Jain and Bari, 2010; Reyes-Chilpa et al., 2006). Among these uses include the treatment of gastrointestinal disorders, leprosy, fevers, and cataracts (Mesía-Vela et al., 2001; Reyes-Chilpa et al., 2006). Numerous *Lonchocarpus* species have been shown to be rich sources of polyphenolic compounds. Polyphenolic compounds are secondary metabolites that meet the demands of plants, although they are not required for normal growth, development, or reproduction (Cseke et al., 2006).

The structural roles in various supporting or protective issues, allelopathic activity, defense-related activities, UV protection, pollinator attraction, and signal molecules involved in interactions between the plant and its environment are among the functions of these secondary metabolites, according to Jaganath and Crozier (2010). Polyphenols are widely distributed in nature, particularly in *Lonchocarpus* plants (Regnault-Roger et al., 2004). They are

found in cereal, tea, coffee, wine, fruits, and vegetables. Their biological properties, such as their anti-inflammatory, anti-ulcer, anti-cancer, antiviral, antifungal, and antioxidant action, make them noteworthy (Hatti et al., 2009). Interest in polyphenolic compounds has grown as a result of these initiatives.

The ability of polyphenolic substances to release electrons, which can subsequently help change the redox state of metal-containing enzymes or interact with the free radicals produced at the enzyme's active site, is the basis for their antioxidant effect. In either scenario, polyphenolic compounds may affect the enzyme activity (Monin et al., 2000). Stilbenes and flavonoids are only two examples of the many structural variants of polyphenolic compounds (Borges-Argáez et al., 2002; Magalhães et al., 2007; Fontenele et al., 2005; Borges-Argáez et al., 2007). They are composed of many families, including aurones, flavanones, pterocarpanes, dibenzoylmethane derivatives, rotenoids, flavanols, flavones, flavans, chalcones, and isoflavonoids (Lima et al., 2009; Magalhães et al., 2007; Cseke et al., 2006; Cassidy et al., 2011). Williams and Grayer (2004) claim that *lonchocarpus* contains a large number of flavonoids with distinctive substitutions. A number of *lonchocarpus* species have flavanones with prenyl and dimethylpyrano group linkages, according to Ingham et al. (1988). Numerous substituents were found to be shared by several of these flavonoids. One such commonly occurring substitution is the prenyl (3-methyl-2-butenyl) group (Morante-Carrie et al., 2024).

Rotenoids with insecticidal and pesticidal properties, such as deguelin and rotenone, are commonly found in this genus (Caboni et al., 2004). Deguelin, a rotenoid, can be injected directly into tumors and is useful in the treatment of cancer (Hebert et al., 2004). Numerous insect species have been shown to be susceptible to the insecticidal effects of rotenone and certain other rotenoids, including deguelin and tephrosin (Fukami and Nakajima, 1971). The roots of *Lonchocarpus* and *Derris* species in South America and Asia, respectively, provide the majority of economically produced rotenone, according to Fukami and Nakajima (1971). The root bark of the native Central American plant *Lonchocarpus guatemalensis* was extracted in methanol to yield the flavanones lupinifolinol, 3-O-methyl-lupinifolinol, and lupinifolin (Magalhães et al., 2007).

Recently, two flavonoids that are novel to *Lonchocarpus araipensis* (Benth.) were identified: furanoflavan and benzofuran-3-ol. These flavonoids have antiallergic, anti-platelet, antioxidant, antiviral, anti-inflammatory, and antitumor effects, according to Lima et al. (2014).

3. PHYTOCHEMISTRY Of *Lonchocarpus* SERIES

3.1 Auronols and Aurones

The majority of *Lonchocarpus* series include aurones and auronols. These are compounds found in fruits and flowers that affect the color of the plants those chemicals are found (Mazziotti et al., (2021); Boumendjel, (2003). Both aurones and auronols, which are closely related, have a five-membered ring. The location of the carbonyl group distinguishes the two types of molecules. Simply, aurones are saturated dihydroaurones. The earliest known auronols were found in *Lonchocarpus obtusus*. Names were assigned to Derriobtusones A and B (Nascimento et al., 1976). Since then, Derriobtusone A, a very rare auronol, has been found in the *Lonchocarpus* species *L. montanus*. In fact, derriobtusone A was one of the substances that were most prevalent in *L. montanus* (Jain and Bari, 2010).

3.2 Chalcones

The lack of a pyran ring distinguishes chalcones from other flavonoids. The pyran ring found in other flavonoids is absent from chalcones (Panche et al., 2016). According to Cseke et al. (2006), they are more conjugated and frequently colored. Their chalcones exhibit antibacterial, antitumoral, anti-inflammatory, cytotoxic, and antioxidant properties against a range of pathogens, including bacteria, viruses, and protozoa, according to Fontenele et al. (2009), De-Andrade et al. (2003), and Fontenele et al. (2005). To date, 29 prenylated chalcones have been isolated from several species of *Lonchocarpus*.

The majority of these were isolated from the seeds and roots of the plants. Nevertheless, it was demonstrated that 3-O-methylabyssinone, the first chalcone isolated from *L. nicou*, is the only chalcone in the genus *Lonchocarpus* that is prenylated on the B-ring (Lawson et al., 2010). It was discovered that *Lonchocarpus* contains dihydrochalcone Erioschalcone B. It was originally documented from tetrahydroflemichapparin-A semisynthesis.

Eriosema glomerata was used to create a unique natural material with antibacterial qualities that was described (Yam-Puc and Pea-Rodriguez, 2009).

3.3 Flavanones, Flavanols, and Flavans

Flavanones are a modification of chalcones. The core structure of flavanols is identical, but the middle ring has an extra hydroxyl group. Although the carbonyl oxygen is changed into a hydroxyl or methoxy group, the fundamental structural properties of flavans are still present (Rosa et al., (2019); Gomes et al., (1981). Only this particular flavanol was found in the species *Lonchocarpus*. The first unique 2,4-dioxygenated flavans of this sort were found in nature when they were isolated from *L. muehlbergianus* (Gomes et al., 1981).

2,4,5,8-tetramethoxyfurano [7,6:2",3"] flavan was found to be the highest abundant flavonoid in the extract. Research on the biosynthesis pathways to flavones and dibenzoylmethanes offers insight into the potential production of these compounds. According to Jain and Bari (2010), a 2-hydroxyflavanone derivative is changed into a 4-hydroxyflavanone derivative, which can subsequently undergo further alkylation. *L. xuul* was used to isolate 4-ethoxy-5-methoxy-6",6"-dimethyl-2H-pyrano-[2",3":7,6]-flavan, another distinct flavan. Since ethoxyl derivatives among flavonoids are rather rare, it is thought that this might not be a true metabolite but rather a byproduct of the isolation process (Borges-Argáez et al., 2000). Since it was the sole ethoxy chemical discovered from *Lonchocarpus*, it is possible that it is just an artifact (Magalhães et al., 2007).

3.4 Isoflavonoids, Flavones and Flavonols in *Lonchocarpus*

Of all the classifications, flavones and flavonols are the most common. It has been discovered that certain flavonols have antibacterial properties (Chagas et al., (2022); Cseke et al., 2006). The pharmacological effects of known flavones include anti-inflammatory, antispasmodic, antinociceptive, anti-allergic, antioxidant, anti-ulcerogenic, antiviral, and antitumoral properties, according to study by Braz et al. (1975). Because ring B is joined to the carbon next to the carbonyl carbon rather than the carbon next to the oxygen, isoflavonoids have a different structure than other flavonoids (Braz et al., 1975).

Previous investigations have identified a number of flavonoids with antioxidant qualities in the leaves, stems, and root barks of *Lonchocarpus xuuul* and *Lonchocarapus yucatenesis* trees on the Yucatan Peninsula (Borques-Argaez et al., 2007). The cytotoxic and antiprotozoal properties of the flavonoids were shown to be moderately active (Borques-Argaez et al., 2007). These flavonoids included 5,4'-dihydroxy-3'-methoxy (6:7), 2',4-dimethoxy-6'-hydroxylonchocarpin, Carpachromene and 2,2 Dimethylpyranoflavone, together with 5,4'-Dimethoxy-(6:7) (Magalhães et al., 2007).

Several flavonoids have been found to have several functions. These include anti-cancer, anti-inflammatory, antibacterial, antiviral, antitumor, cytotoxic, and antiprotozoal qualities, as well as the potential to reduce the risk of cardiovascular disease. This range of activities may be connected in some way to the antioxidant qualities of flavonoids (Borges-Argáez et al., 2007; Jain and Bari, 2010; Blatt et al., 2002).

Cseke et al. (2006) claim that isoflavonoids have antifungal, estrogenic, insecticidal, and anticancer properties. Isoflavonoids include both isoflavones and isoflavans. In contrast to isoflavans, which are comparable to flavans in that they do not have the double bond in ring C or the carbonyl oxygen, isoflavones share these characteristics with flavones. *Lonchocarpus* contained a relatively high amount of flavones. Numerous isolated flavones from different species of *Lonchocarpus* were found. Many of the compounds were found in many species. Quercetin was the only flavonol found in the *Lonchocarpus* species. Numerous plants and foods contain this well-known flavonol, which has a broad spectrum of biological activities (Xueqin et al., 2011; Magalhães et al., 2007).

3.5 Pterocarpan in *Lonchocarpus*

According to Gomes et al. (1981), Perez et al., (2023), pterocarpan is made from isoflavones, where the carbonyl oxygen now forms a part of a five-membered ring. These substances have not been detected as often in *Lonchocarpus* species. Additionally, they were less prevalent among the various species. When they were first found, each species usually had at least two of them. Three pterocarpanes were identified in *Lonchocarpus montanus*, but medicarpin was the only one detected in many *Lonchocarpus* species. Philenopteran and 9-O-

Methylphenlopteran were the only two pterocarpanes found in *L. laxiflorus* (Magalhães et al., 2007).

3.6 Rotenoids in *Lonchocarpus*

Based on the isoflavonoid skeleton, which includes rotenoids, *Lonchocarpus* generates a wide range of secondary polyphenolic metabolites (Magalhães et al., 2007; Lawson et al., 2010). *Lonchocarpus* is recognized for its high amounts of rotenoid chemicals, even though it has been proven to contain a range of flavonoids. Because they contain rotenoid compounds, many species of *Lonchocarpus* are toxic to fish and insects (Mesía-Vela et al., 2001; Reyes-Chilpa et al., 2006).

Although they are not as prevalent as the other flavonoids, rotenoids are found in *Lonchocarpus* (Magalhães et al., 2007). Interest has increased in trying to comprehend the composition and biological roles of plants that contain rotenoid. This is because, in addition to their use as agrochemicals, they may also be useful as anti-carcinogenesis agents (Fang et al., 1999). In addition to rotenone itself, many other derivatives have been found, including epoxides of particular interest. It is believed that the two rotenone epoxides found in *L. nicou* roots are the first to be derived from plants (Lawson et al., 2010).

Another class of rotenone and deguelin derivatives has been discovered, but this time they come from both *L. nicou* and *L. urucu*. The skeleton of these new rotenoid-like compounds is unique, featuring chromone rings on the C and D and 1,5-benzodioxepin rings on the A and B. These specific compounds were present in a plant extract for the first time. They were proven to be authentic natural compounds rather than extraction byproducts (Fang et al., 1999).

3.7 Stilbenes in *Lonchocarpus*

Stilbenes, which are made up of two benzene rings connected by an ethene, are the most fundamental kind of polyphenolics. Stilbenes have been shown to have substituents that are similar to those found in flavonoids. These include methoxy, hydroxyl, and prenyl groups. Hydroxystilbenes have been identified in a wide variety of plants. They are usually hydroxylated at 3, 5, 3, 4', 5, 2, 3, 5, or 3, 3', 5, according to Xueqin et al. (2011). These patterns may also contain prenyl and methyl groups. Prenylated stilbenes are commonly found in *lonchocarpus* (Magalhães et al., 2007). Numerous substances

include at least one prenyl group. These consist of 3,5-dimethoxy-4-prenylstilbene, 3,4',5-trimethoxy-4-prenylstilbene, longistylins A through D, and chiricanins A through E (Magalhães et al., 2007).

3.8 Coumaronochromones, 3-Phenylcoumarins and Acids in *Lonchocarpus*

Blatt et al. (2002) state that isoflavones that have undergone additional C-2 oxidation are 3-phenylcoumarins. They are less common than some of the other flavonoids because they are the only molecules in the flavonoid class that contain ester and that can undergo C-2 oxidation. Of these specific compounds, only three were found in species of *Lonchocarpus*. However, it may be important for these types of compounds that two were found to be from the same plant (Shah and Smith, 2020; Blatt et al., 2002).

According to Lawson et al. (2010), the B-ring of coumaronochromones, a type of rotenoid, has

been compressed into a five-membered ring. The first examples of a prenylated benzoic acid were found in *Lonchocarpus nicou*. The asteraceae family has been found to possess prenylated phenolic acids, despite their rarity in nature (Xueqin et al., 2011).

Many *Lonchocarpus* species have been used as diuretics, laxatives, stomachics, parasiticides, and to treat tumors, eruptions, scurvy, backaches, convulsions, and wounds (Emmanuel et al., (2020); Mesía-Vela et al., (2001); Reyes-Chilpa et al., (1995). Because polyphenolic chemicals are poisonous and/or antifeedant to insects that are crop pests, several species of *Lonchocarpus* are protected from insect assault (Simmonds et al., 1990). Borges-Argáez et al. (2007) state that *lonchocarpus* is well known for its insecticidal and pesticidal qualities.

According to Dos-Santos et al. (2009), *Lonchocarpus* species have also been used as anthelmintics, cough suppressants, antidiarrheal, antileprosy drugs, and fish poisons. In traditional

Table 1. Some *Lonchocarpus* species and their ethnomedical uses

<i>Lonchocarpus violaceus</i> (Balche) Root bark Stem bark	Stress reduction, an alternative to alcohol and an intoxicating substance, an anticancer, a mosquito repellent, fish poison, and an insecticide.	Udeani et al., 1997; Correa, 1984; Sule et al., 2022.
<i>Lonchocarpus eriocalyx</i>	Fever, malaria, a skin condition, thrush, and ringworms ringworm treatment, insect repellent, and headache alleviation	Kareru et al., 2007 Yenesew et al., 2004
<i>Lonchocarpus costaricensis</i>	Insecticide, Treatment of thrush	Caboni et al., 2004
<i>Lonchocarpus araripensis</i>	Rheumatism, diabetes, skin diseases and arthritis.	Julianeli et al., 2011
<i>Lonchocarpus urucu</i> (Barbasco) Root bark Stem bark	Anti-HIV, anti-insect, anti-pest, anti-headache, and treatment for skin conditions.	Udeani et al., 1997 Correa, 1984
<i>Lonchocarpus oliganthus</i>	Acts as Insecticide	Xueqin et al., 2011
<i>Lonchocarpus cyanescens</i>	Ulcers, Antioxidant, arthritis	Samwel, et al., 2014
<i>Lonchocarpus laxiflorus</i>	Back pain, rashes, ulcers, intestinal worms, mental disease, headaches, and back discomfort.	Neuwinger, 1996 Okello and Ssegawa, 2007
<i>Lonchocarapus Yucatenensis</i>	Headache and Fever	Borges- Argaez et al., 2007
<i>Lonchocarpus chiricanus</i>	Insecticide	Jean-Robert et al., 2001
<i>Lonchocarpus xuuul</i>	headache and fever	Borges- Argaez et al., 2007
<i>Lonchocarpus guatemalensis</i>	Fever, headache	Xueqin et al., 2011
<i>Lonchocarpus castilloi</i>	Antifungal and Anti-termite	Alavez-Solano et al., 2000

Nigerian recipes, the medicinal herb *Lonchocarpus cyanescens* (LC) is sometimes used in combination with other components to cure mental illnesses. *Lonchocarpus cyanescens* is the primary component of the Nigerian psychosis treatment formula (Sonibare et al., 2012).

3.9 *Lonchocarpus eriocalyx* (Harms)

Lonchocarpus eriocalyx is a member of the *Fabaceae* family, which generates a wide range of flavonoids with insecticidal and other antimicrobial properties (Roy et al., (2022); Ceres et al., 1981). A previous study found that a crude extract from the root bark of *L. eriocalyx* was effective against *P. falciparum* strains that were resistant to chloroquine (D6) and susceptible to it (W2) (Tuwei, 2006). Furthermore, this extract demonstrated larvicidal activity against larvae of *A. aegypti* mosquitoes. The single molecule identified from *L. eriocalyx* was lupeol, a lupane-type triterpene with strong antiparasitic activity (Yenesew et al., 2004; Tuwei, 2006).

3.10 *Lonchocarpus cyanescens* (LC)

The aerial parts of *Lonchocarpus cyanescens* yield an indigo that has been used as a valuable dye for garments in West Africa from ancient times (adire/gara), according to Perkin and Thomas (1909), Dorcas and Ibrahim, (2013). For court penalties, ceremonial outfits, burial garments, bridal dowries, and gifts for important guests, chiefs typically use woven fabric known as "country cloth" dyed with "gara" (both natural and chemical dyes). This dye comes from the "gara" leaf that was recovered from LC (Dorcas and Ibrahim, 2013; Cardon, 2003; Spencer, 1996; Polakoff, 1980). "Gara" is a product that is sold extensively in the industrial sector; dried LC is exported to Europe, the United States, the United Kingdom, and many other African nations from Liberia and other West African nations (Cardon, 2003; Spencer, 1996; Polakoff, 1980).

Historically, the plant has been used for medicinal purposes. The leaves are served alongside couscous in Senegal. In Ghana, it is thought that LC roots work better than leaves for curing ulcers and other skin conditions (Boakye-Yiadom et al., 2021). The leaves and roots of this plant have also been used to treat leprosy in Guinea-Bissau and Sierra Leone. Leaf material can be used to make laxatives (Spencer, 1996). In Benin, leaf sap is used to treat gastrointestinal issues and diarrhea. In addition to being used as

an aphrodisiac, a decoction of leafy twigs and roots is administered to women during or after childbirth. In Nigeria, this mixture is used to treat rheumatism, diarrhea, and venereal diseases. Ground root is used to alleviate yawns, and water with powdered root is used to wash sores (Irvine, 1961; Spencer, 1996; Jukema et al., 1991; Iwu et al., 2014). The tropical plant LC has long been used by fisherman in Nigeria's Akwa Ibom State to scare and capture fish. LC is a shrub belonging to the *Dalbergiaceae* tribe of the *Leguminosae* natural order, which provides a habitat for twining (Iwu et al., 2006).

It is listed among the medicinal plants of Akwa Ibom State (Ajibesin et al., 2008). LC is well-known since it is used to create dye. The blue indigo dye, which is used to color textiles and other materials, is fermented from the indican-containing early sprouts and leaves (Ajibesin et al., 2008). Indigotin, a component of indigo dye, is produced by indoxyl, which is present in LC leaves. Oleanane derivatives and glycyrrhetic acid are responsible for the anti-inflammatory properties and therapy of peptic ulcers observed in LC (Cardon and Jarsen 2005; Iwu et al., 2006; Ogbeide et al., 1991; Promsattha et al., 1987). According to Cardon and Jarsen (2005), Iwu et al. (2006), and Ndukwe et al. (2004), triterpenes exhibit anti-arthritis qualities. These natural elements also contribute to the flavoring's taste effects (Ijichi and Seizo, 2005). It has been demonstrated that the bioactivity effects of *Lonchocarpus cyanescens* are ulcer-relieving, anti-inflammatory, and anti-arthritis (Iwu et al., 2006).

A phyto-constituents investigation revealed that *L. cyanescens* bark aqueous extracts contained alkaloids, flavonoids, saponins, phytates, glycosides, and oxalates. Alkaloids, flavonoids, and glycosides were just slightly present, whereas oxalates were widely distributed. Phytates and saponins were also detected in trace levels (Xueqin et al., 2011).

3.11 *Lonchocarpus sericeus* (Poir.)

Often referred to as Senegal lilac or Cube root, *Lonchocarpus sericeus* *Papilionaceae* is a dry deciduous tree that grows natively in Ghana and Nigeria. It can grow up to 16 meters in height. In traditional African medicine, the bark is mostly used as a laxative and stomachic, especially for young children (Paul and Olajide, 2023). Additionally, it is frequently used to treat back pain and convulsions (Burkill, 1995). Its

blossoms, which have many racemes of purple flowers dangling from them, make it ideal for presentations. The flowers have a unique, almost vanilla-like fragrance. It is commonly planted as a shade tree in communities and in gardens. Kojis et al., (2004) and Adewuyi et al., (2012) stated that the wood is translucent yellow with sporadic marbling and has an olive-green heartwood. The leaves are used for universal healing, laxatives, and fish poisons in Nigeria, while the bark is used to treat body aches, arthritis, rheumatism, cutaneous and subcutaneous parasite infections, malnourishment, debility, paralysis, and convulsions. The roots are used to treat leprosy. The seeds and berries are used as insect repellents called andarachnicides (Burkill, 1985). Additionally, it is utilized as body cream in cosmetics.

3.12 *Lonchocarpus bussei* (L. Laxiflorus)

A plant species called *Lonchocarpus bussei*, a legume belonging to the *Fabaceae* family, is found in Nigeria. African snowdrop tree and West African snowdrop tree are common names for it (Paul and Olajide, 2023). *Lonchocarpus bussei* is a deciduous tree that can grow up to 30 meters in height. It has smooth gray bark and a straight trunk. The leaves are pinnate, compound, and arranged alternately along the stem. Numerous leaflets, often numbering five to eleven, make up each leaf. The tree produces panicles, which are small, clustered blooms that range in color from white to pale yellow. During the blossoming season, the blossoms are abundant and have a unique fragrance (Moore and Bradley, 2022). The flat, woody pods that *Lonchocarpus bussei* produces are typically 7 to 20 centimeters long. The pods are initially green, but as they mature, they turn brown. Each pod contains several seeds. *Lonchocarpus bussei* is widespread in several countries in West and Central Africa, including Nigeria, Cameroon, Gabon, and the Democratic Republic of the Congo (Afolabia et al., 2019). Strong and highly valued, *lonchocarpus bussei* wood is used in building. There are other traditional uses for the tree as well. In Nigeria, for example, the bark is sometimes used as a medicine to treat particular ailments, and the leaves are also utilized in traditional remedies (Paul and Olajide, 2023).

4. CONCLUSION

Polyphenolic compounds are common in *lonchocarpus* species. Numerous types of polyphenolic chemicals, including chalcones,

flavans, aurones, dibenzoylmethane derivatives, rotenoids, pterocarpans, flavanones, flavanols, flavones, isoflavones, and flavonols, are abundant in *Lonchocarpus*, according to phytochemical research. Although a thorough examination has not yet been conducted, the plant genus *Lonchocarpus* has a number of chemicals with a wide range of varied properties. In Africa, *lonchocarpus sericeus* is used to promote appetite, treat stomach issues, treat epilepsy, and act as a laxative. *Lonchocarpus* has several traditional applications. Occasionally, traditional medicine uses the bark and leaves to treat a variety of ailments. Currently under discussion are three species of *Lonchocarpus*: *Lonchocarpus cyanescens*, *Lonchocarpus bussei*, and *Lonchocarpus sericeus*. *Lonchocarpus* has not gotten much study attention, even though several of its species contained tannins, flavonoids, rotenoids, and isoflavonoids. These phytochemicals support the traditional applications and therapeutic qualities of *Lonchocarpus* species in a number of indigenous medical systems. To completely understand the phytochemical profile of the *Lonchocarpus* series, as well as its unique ingredients, methods of action, and any adverse effects on people, more investigation and study are therefore required.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Abdullahi, S. M., Musa, A. M., Abdullahi, M. I., Sule, M. I., & Sani, Y. M. (2013). Isolation of Lupeol from the stem-bark of *Lonchocarpus sericeus* (Papilionaceae). *Scholars Academic Journal of Biosciences*, 3, 18-19.
- Adewuyi, R. A., Oderinde, B. V. S. K., Rao, R. B. N., & Prasad, R. B. N. (2012). Chemical composition and molecular speciation of the triacylglycerol of the oils of *Lonchocarpus sericeus* and *Lonchocarpus*

- cyanescens*. *Natural Product Research*, 26(20), 1954-1956.
- Afolabi, C. G., Ogunsanya, O. M., & Lawal, O. I. (2019). Evaluation of some African yam bean (*Sphenostylis stenocarpa* [Hochst. Ex A. Rich]) accessions for resistance to flower bud and pod rot diseases. *Current Plant Biology*, 20, 100126.
- Ajibesin, K. K., Ekpo, B. A., Bala, D. N., Essien, E. E., & Adesanya, S. A. (2008). Ethnobotanical survey of Akwa Ibom State of Nigeria. *Journal of Ethnopharmacology*, 115, 387-408.
- Alavez-Solano, D., Reyes-Chilpa, R., Jimenez-Estrada, M., Gomez-Garibay, F., Chavez-Urbe, I., & Sousa-Sanchez, M. (2000). Flavanones and 3-hydroxyflavanones from *Lonchocarpus* *oaxacensis*. *Phytochemistry*, 55, 953-957.
- Bentjee, H. (1994). *Kenyan Trees, Shrubs and Lianas*. National Museums of Kenya, Nairobi, 102-110, 296.
- Blatt, C. T. T., Chávez, D., Chai, H., Graham, J. G., Cabieses, F., Farnsworth, N. R., Cordell, G. A., Pezzuto, J. M., & Kinghorn, A. D. (2002). Cytotoxic flavonoids from the stem bark of *Lonchocarpus* aff. *fluvialis*. *Phytotherapy Research*, 16, 320-325.
- Boakye-Yiadom, M., Kumadoh, D., Adase, E., & Woode, E. (2021). Medicinal plants with prospective benefits in the management of peptic ulcer diseases in Ghana. *Biomed Research International*, 5574041. <https://doi.org/10.1155/2021/5574041>
- Borges-Argáez, R., Balnburry, L., Flowers, A., Giménez-Turba, A., Ruiz, G., Waterman, P. G., & Peña-Rodríguez, L. M. (2007). Cytotoxic and antiprotozoal activity of flavonoids from *Lonchocarpus* spp. *Phytomedicine*, 14, 530-533.
- Borges-Argáez, R., Peña-Rodríguez, L. M., & Waterman, P. G. (2000). Flavonoids from the stem bark of *Lonchocarpus xuul*. *Phytochemistry*, 54, 611-614.
- Borges-Argáez, R., Peña-Rodríguez, L. M., & Waterman, P. G. (2002). Flavonoids from two *Lonchocarpus* species of the Yucatán Peninsula. *Phytochemistry*, 60, 533-540.
- Borges-Argáez, R., Vela-Catzín, T., Yam-Puc, A., Chan-Bacab, M. J., Moo-Puc, R. E., & Cáceres-Farfán, M. (2009). Antiprotozoal and cytotoxic studies on some isocordoin derivatives. *Planta Medica*, 75, 1336-1338.
- Boumendjel, A. (2003). Aurones: A subclass of flavones with promising biological potential. *Current Medicinal Chemistry*, 10(23), 2621-2630.
- Braz Filho, R., Gottlieb, O. R., Mourão, A. P., Da Rocha, A. I., & Oliveira, F. S. (1975). Flavonoids from *Derris* species. *Phytochemistry*, 14, 1454-1456.
- Burkill, H. M. (1985). Entry for *Lasiurus hirsutus* (Forssk.) Boiss [Family Poaceae]. In H. M. Burkill (Ed.), *The Useful Plants of West Tropical Africa* (2nd ed.). Royal Botanical Garden, Kew, UK.
- Burkill, H. M. (1995). *The Useful Plants of West Tropical Africa* (2nd ed.). Vol. 3. Royal Botanic Gardens, Kew. United Kingdom.
- Caboni, P., Sherer, T., Zhang, B., Taylor, G., Grenamyre, J., Timothy, C., & John, E. (2004). Rotenone, deguelin, and their metabolites. *Chemical Research in Toxicology*, 17(11), 1540-1548.
- Cardon, D. (2003). *Le Monde des Teintures Naturelles*. Belin, Paris.
- Cardon, D., & Jarsen, P. C. M. (Eds.). (2005). *Philenoptera cyanescens* (Schumach. & Thonn.) Roberty. In P. C. M. Jarsen & D. Cardon (Eds.), *PROTA 3: Dyes and Tannins/Colorants et Tannins* (pp. 194-196). PROTA, Wageningen.
- Cassidy, E., Caittli, N., & William, S. (2011). *Lonchocarpus* polyphenolics and their biological activities. *The Natural Products Journal*, 1(2), 75-104.
- Ceres, M., Gottlieb, R., Giovanno, B., Marini, B., France, D., & Roger, P. (1981). Seven flavonoids from some Iranian angiosperms. *Biochemical Systematics and Ecology*, 9(2-3), 129-147.
- Chaachouay, N., & Zidane, L. (2024). Plant-derived natural products: A source for drug discovery and development. *Drugs and Drug Candidates*, 3(1), 184-207. <https://doi.org/10.3390/ddc3010011>
- Chagas, M. D. S. S., Behrens, M. D., Moragas-Tellis, C. J., Penedo, G. X. M., Silva, A. R., & Gonçalves-de-Albuquerque, C. F. (2022). Flavonols and flavones as potential anti-inflammatory, antioxidant, and antibacterial compounds. *Oxidative Medicine and Cellular Longevity*, 2022, 9966750. <https://doi.org/10.1155/2022/9966750>
- Correa, M. P. (1984). *Dicionário das plantas úteis do Brasil e das exóticas cultivadas*. IBDF, Ministério da Agricultura, Rio de Janeiro.
- Cseke, L. J., Kirakosyan, A., Kaufman, P. B., Warber, S., Duke, J. A., & Briemann, H. L. (2006). *Natural products from plants* (2nd ed.). CRC Press, Boca Raton, FL.

- De-Andrade Cunha, G. M., Fontenele, J. B., Nobre Júnior, H. V., de Sousa, F. C. M., Silveira, E. R., Nogueira, N. A. P., de Moraes, M. O., Viana, G. S. B., & Costa-Lotufo, L. V. (2003). Cytotoxic activity of chalcones isolated from *Lonchocarpus sericeus* (Pocr.) Kunth. *Phytotherapy Research*, 17, 155-159.
- Dos-Santos, D. A. P., Braga, P. A. C., da Silva, M. F. G. F., Fernandes, J. B., Vieira, P. C., Magalhães, A. F., Magalhães, E. G., Marsaioli, A. J., Moraes, V. R. S., Rattray, L., & Croft, S. L. (2009). Anti-African trypanocidal and antimalarial activity of natural flavonoids, dibenzoylmethanes, and synthetic analogues. *Journal of Pharmaceutical and Pharmacological Sciences*, 61, 257-266.
- Emmanuel, U., Onyekachi, J., & Uchenna, O. (2020). Phytochemical, anti-inflammatory and analgesic properties of stem-bark extract and fractions of *Lonchocarpus sericeus* Poir. (Papilionaceae) in albino mice. *Avicenna Journal of Phytomedicine*, 10(6), 584-593. <https://doi.org/10.22038/ajp.2020.48207.2383>
- Estrada-Castillón, E., Villarreal-Quintanilla, J. Á., Cuéllar-Rodríguez, G., Torres-Colín, L., Encina-Domínguez, J. A., Sánchez-Salas, J., Muro-Pérez, G., González-Cuéllar, D. A., Galván-García, O. M., Rubio-Pequeño, L. G., & Mora-Olivo, A. (2024). The Fabaceae in northeastern Mexico (Subfamilies Caesalpinioideae, Excluding Tribe Mimoseae, Cercidoideae, and Detarioideae). *Plants*, 13(17), 2477. <https://doi.org/10.3390/plants13172477>
- Fang, N., & Casida, J. E. (1999). New bioactive flavonoids and stilbenes in cube resin insecticide. *Journal of Natural Products*, 62(2), 205-210. <https://doi.org/10.1021/np990031g>
- Farnsworth, N. R., Akerele, O., Bingel, A. S., Soejarto, D. D., & Guo, Z. (1985). Medicinal plants in therapy. *Bulletin of the World Health Organization*, 63, 965-981. [https://doi.org/10.1016/0378-8741\(87\)90016-X](https://doi.org/10.1016/0378-8741(87)90016-X)
- Fontenele, J. B., Leal, L. K. A. M., Félix, F. H. C., Silveira, E. R., & Viana, G. S. B. (2009). Studies on the anti-oedematogenic properties of a fraction rich in lonchocarpin and derricin isolated from *Lonchocarpus sericeus*. *Natural Product Research*, 23(18), 1677-1688. <https://doi.org/10.1080/14786410802252752>
- Fontenele, J. B., Leal, L. K. A. M., Ferreira, M. A. D., Silveira, E. R., & Viana, G. S. B. (2005). Antiplatelet effect of lonchocarpin and derricin isolated from *Lonchocarpus sericeus*. *Pharmaceutical Biology*, 43(8), 726-731. <https://doi.org/10.1080/13880200500397661>
- Fukami, H., & Nakajima, M. (1971). *Rotenone and rotenoids: Naturally occurring insecticides*. Marcel Dekker Inc., New York.
- Gomes, C. M. R., Gottlieb, O. R., Bettolo, G. B. M., Delle Monache, F., & Polhill, R. M. (1981). Systematic significance of flavonoids in *Derris* and *Lonchocarpus*. *Biochemical Systematics and Ecology*, 9(2/3), 129-147. [https://doi.org/10.1016/0305-1978\(81\)90047-X](https://doi.org/10.1016/0305-1978(81)90047-X)
- Haegnauer, R., & Grayer-Barkmeijer, R. (1993). Relevance of seed polysaccharides and flavonoids for the classification of the leguminosae: A chemotaxonomic approach. *Phytochemistry*, 34, 3-16. [https://doi.org/10.1016/0031-9422\(93\)80244-N](https://doi.org/10.1016/0031-9422(93)80244-N)
- Hatti, K. S., Diwakar, L., Rao, G. V., Kush, A., & Reddy, G. C. (2009). Abyssinones and related flavonoids as potential steroidogenesis modulators. *Bioinformation*, 3(9), 399-402. <https://doi.org/10.6026/9732063000399>
- Hebert, K., Paul, D. N., Penton, L., Erin, H., Burns, K., Janzen, M., Daniel, H., Hallawachs, W., Beier, J. C., Perkins, P. V., & Onyango, F. K. (2004). Ten species in one: DNA barcoding reveals cryptic species in the semi-tropical skipper butterfly *Astraptes fulgerator*. *Proceedings of the National Academy of Sciences*, 101(41), 14812-14817. <https://doi.org/10.1073/pnas.0403992101>
- Ijichi, S., & Seizo, T. (2005). Molecular design of sweet tasting compounds based on β -amino-3 β -deoxy-18 β -glycyrrhetinic acid: Amido functionality eliciting tremendous sweetness. *Chemistry Letters*, 34, 356-357. <https://doi.org/10.1246/cl.2005.356>
- Ingham, J. L., Tahara, S., & Dziedzic, S. Z. (1988). Major flavanones from *Lonchocarpus guatemalensis*. *Zeitschrift für Naturforschung C*, 43, 818-822. <https://doi.org/10.1515/znc-1988-0611>

- Irvine, F. R. (1961). *Woody plants of Ghana, with special reference to their uses*. Oxford University Press, London.
- Iwu, M. M. (2006). Comparative estimation of glycyrrhetic acid content of *Lonchocarpus cyanescens* and *Glycyrrhiza* roots. *Journal of Pharmaceutical Sciences*, 71(12), 1412-1413. <https://doi.org/10.1002/jps.2600711227>
- Iwu, M. M. (2014). Pharmacognostical profile of selected medicinal plants. In *Handbook of African Medicinal Plants* (pp. 251-252). CRC Press, Florida.
- Jaganath, I. B., & Crozier, A. (2010). Dietary flavonoids and phenolic compounds. In C. G. Fraga (Ed.), *Plant phenolics and human health* (p. 2). John Wiley & Sons, Inc.
- Jain, P. S., & Bari, S. B. (2010). Isolation of Lupeol, Stigmasterol and Campesterol from petroleum ether extract of woody stem-bark of *Wightia tinctoria*. *Asian Journal of Plant Sciences*, 9(3), 163-167. <https://doi.org/10.3923/ajps.2010.163.167>
- Jean-Robert, I., Andrew, M., Mahabir, P. G., & Kurt, H. (2001). Five new prenylated stilbenes from the root of *Lonchocarpus chiricanus*. *Journal of Natural Products*, 64(6), 710-715. <https://doi.org/10.1021/np0100446>
- Jukema, J. N., Wulijarni-Soetjipto, R. H., Lemmens, J. W., & Hildebrand, J. W. (1991). Minor dye and tannin-producing plants. In R. H. Lemmens & N. Wulijarni-Soetjipto (Eds.), *Plant Resources of South-East Asia No 3: Dye and Tannin Producing Plants* (pp. 132-142). Pudoc.
- Julianeli, T., Jackson, R. G., Kelly, S. A., Sílvia, S. C., Celso, S. C., & José Maria, B. F. (2011). Selective spasmolytic effect of a new furanoflavoquinone derivative from diplotropin on guinea-pig trachea. *Journal of Chemical and Pharmaceutical Research*, 3(1), 249-258.
- Kareru, P. G., Kenji, G. M., Gachanja, A. N., Keriko, J. M., & Mungai, G. (2007). Traditional medicines among the Embu and Mbeere people of Kenya. *African Journal of Traditional, Complementary and Alternative Medicines*, 4(1), 75-86. <https://doi.org/10.4314/ajtcam.v4i1.31385>
- Kojs, P., Włoch, W., & Rusin, A. (2004). Rearrangement of cells in storeyed cambium of *Lonchocarpus sericeus* (Poir.) DC connected with formation of interlocked grain in the xylem. *Trees*, 18, 136-144. <https://doi.org/10.1007/s00468-003-0307-6>
- Kokwaro, J. O. (2009). *Medicinal plants of East Africa* (3rd ed., pp. 134-137). University of Nairobi Press.
- Lawson, M. A., Kaouadji, M., & Chulia, A. J. (2010). A single chalcone and additional rotenoids from *Lonchocarpus nicou*. *Tetrahedron Letters*, 51, 6116-6119. <https://doi.org/10.1016/j.tetlet.2010.08.133>
- Lima, A. F., Ferreira, D. A., & Monte, F. J. (2014). Flavonoids from *Lonchocarpus araripensis* (Leguminosae): Identification and total ¹H and ¹³C resonance assignment. *American International Journal of Contemporary Research*, 4(2), 18-23.
- Lima, A. F., Mileo, P. G. M., Andrade-Neto, M., Braz-Filho, R., Silveira, E. R., & Pessoa, O. D. L. (2009). ¹H and ¹³C NMR assignments of new methoxylated furanoflavonoids from *Lonchocarpus araripensis*. *Magnetic Resonance in Chemistry*, 47, 165-168. <https://doi.org/10.1002/mrc.2422>
- Magalhães, A. F., Tozzi, A. M. G. A., Magalhães, E. G., Sannomiya, M., Soriano, M. D. P. C., & Perez, M. A. F. (2007). Flavonoids of *Lonchocarpus montanus* A.M.G. Azevedo and biological activity. *Anais da Academia Brasileira de Ciências*, 79(3), 351-367. <https://doi.org/10.1590/S0001-37652007000300002>
- Mannetje, L. (2002). *Alysicarpus ovalifolius* (Schumacher): J. Leonard record from Protobase. In L. P. A. Oyen & R. H. J. Lemmens (Eds.), *Plant Resources of Tropical Africa* (pp. 325-416). Bulletin du Jardin Botanique de l'État à Bruxelles, 32(4).
- Mazziotti, I., Petrarolo, G., & La Motta, C. (2021). Aurones: A golden resource for active compounds. *Molecules*, 27(1), 2. <https://doi.org/10.3390/molecules27010002>
- Mesía-Vela, S., Sánchez, R. I., Estrada-Muñiz, E., Alavez-Solano, D., Torres-Sosa, C., Jiménez-Estrada, M., Reyes-Chilpa, R., & Kauffman, F. C. (2001). Natural products isolated from Mexican medicinal plants: Novel inhibitors of sulfotransferases, SULT1A1 and SULT2A1. *Phytomedicine*, 8(6), 481-488. [https://doi.org/10.1016/S0944-7113\(01\)80028-5](https://doi.org/10.1016/S0944-7113(01)80028-5)
- Monin, H., Guo, Q., & Packer, L. (2000). Enzyme inhibition and protein-binding action of the procyanidin-rich French maritime pine bark extract, pycnogenol: Effect on xanthine

- oxidase. *Journal of Agricultural and Food Chemistry*, 48, 5630-5639. <https://doi.org/10.1021/jf000295a>
- Moore, K. A., & Bradley, L. K. (Eds.). (2022). *North Carolina Extension Gardener Handbook* (2nd ed.). NC State Extension. <http://content.ces.ncsu.edu/3-botany>
- Morante-Carriel, J., Živković, S., Nájera, H., Sellés-Marchart, S., Martínez-Márquez, A., Martínez-Esteso, M. J., Obrebska, A., Samper-Herrero, A., & Bru-Martínez, R. (2024). Prenylated flavonoids of the *Moraceae* family: A comprehensive review of their biological activities. *Plants*, 13(9), 1211. <https://doi.org/10.3390/plants13091211>
- Moronkola, D. O., & Oladosu, I. A. (2013). Chemical compositions of *Lonchocarpus cyanescens* Benth. (Fabaceae)—Case study of its volatile oils, and two triterpenoids. *American Journal of Plant Sciences*, 4, 1653-1659. <https://doi.org/10.4236/ajps.2013.48199>
- Moronkola, D. O., Adediji, A. A., Oyewole, I. O., Ogunwande, I. A., & Oladosu, I. A. (2009). Survey and accounts of anti-malaria ethnomedicinal herbs used in South-Western Nigeria and vector repellent activities of their volatile oils. *The African Network for Drugs and Diagnostics (ANDI), 2nd Stakeholders Meeting & Conference*, Medical Research Council (MRC), Cape Town, 4-7 October 2009.
- Nascimento, M. C., Vasconcellos Dias, R. L., & Mors, W. B. (1976). Flavonoids of *Derris obtusa*: Aurones and auronols. *Phytochemistry*, 15, 1554-1558. [https://doi.org/10.1016/S0031-9422\(00\)85030-9](https://doi.org/10.1016/S0031-9422(00)85030-9)
- Ndukwe, K. C., Lamikanra, O., & Okeke, I. N. (2004). Antibacterial activity in plants used as chewing sticks in Africa. *Prous Science Drugs of the Future*, 29(12), 1221. <https://doi.org/10.1358/dof.2004.029.12.853766>
- Neuwinger, D. (1996). *African ethnobotany: Poisons and drugs: Chemistry, pharmacology, toxicology* (pp. 682-683). CRC Press.
- Okello, J., & Ssegawa, P. (2007). Medicinal plants used by communities of Ngai Subcounty, Apac District, Northern Uganda. *African Journal of Ecology*, 45(1), 76-83. <https://doi.org/10.1111/j.1365-2028.2006.00675.x>
- Panche, A. N., Diwan, A. D., & Chandra, S. R. (2016). Flavonoids: An overview. *Journal of Nutritional Science*, 5, e47. <https://doi.org/10.1017/jns.2016.41>
- Paul, K., & Olajide, K. (2023). Lesser known African indigenous tree and fruit plants: Recent evidence from literatures and regular cultivation culture. IntechOpen. <https://doi.org/10.5772/intechopen.104890>
- Perez Rojo, F., Pillow, J. J., & Kaur, P. (2023). Bioprospecting microbes and enzymes for the production of pterocarpan and coumestans. *Frontiers in Bioengineering and Biotechnology*, 11, 1154779. <https://doi.org/10.3389/fbioe.2023.1154779>
- Perkin, A. G., & Thomas, F. (1909). RSC publishing paper XC. Indican, Part II. *Journal of the Chemical Society, Transactions*, 95, 793-807. <https://doi.org/10.1039/ct9099500793>
- Polakoff, C. (1980). *Into indigo—African textiles and dyeing techniques* (p. 243). Anchor Books.
- Regnault-Roger, C., Ribodeau, M., Hamraoui, A., Bareau, I., Blanchard, P., Gil-Munoz, M. I., & Barberan, F. T. (2004). Polyphenolic compounds of Mediterranean Lamiaceae and investigation of orientational effects on *Acanthoscelides obtectus* (Say). *Journal of Stored Products Research*, 40, 395-408. <https://doi.org/10.1016/j.jspr.2003.07.003>
- Reyes-Chilpa, R., Baggio, C. H., Alavez-Solano, D., Estrada-Muñiz, E., Kauffman, F. C., Sánchez, R. I., & Mesia-Vela, S. (2006). Inhibition of gastric H⁺,K⁺-ATPase activity by flavonoids, coumarins, and xanthenes isolated from Mexican medicinal plants. *Journal of Ethnopharmacology*, 105, 167-172. <https://doi.org/10.1016/j.jep.2005.08.039>
- Reyes-Chilpa, R., Viveros-Rodríguez, N., Gómez-Garibay, F., & Alavez-Solano, D. (1995). Antitermitic activity of *Lonchocarpus castilloi* flavonoids and heartwood extracts. *Journal of Chemical Ecology*, 21(4), 455-463. <https://doi.org/10.1007/BF02029173>
- Rosa, G. P., Seca, A. M. L., Barreto, M. D. C., Silva, A. M. S., & Pinto, D. C. G. A. (2019). Chalcones and flavanones bearing hydroxyl and/or methoxyl groups: Synthesis and biological assessments. *Applied Sciences*, 9(14), 2846. <https://doi.org/10.3390/app9142846>
- Roy, A., Khan, A., Ahmad, I., Alghamdi, S., Rajab, B. S., Babalghith, A. O., Alshahrani, M. Y., Islam, S., & Islam, M. R. (2022). Flavonoids: A bioactive compound from medicinal plants and its therapeutic

- applications. *Biomed Research International*, 5445291. <https://doi.org/10.1155/2022/5445291>
- Samwel, B., Adigun, O., & Adaramoye, O. (2014). Bioassay-guided investigation of *Lonchocarpus cyanescens* Benth leaf extract for antioxidant activities. *African Journal of Biotechnology*, 13(22), 2240-2247. <https://doi.org/10.5897/AJB2014.13935>
- Schrire, B. D., Lewis, G. P., & Lavin, M. (2005). Biogeography of the Leguminosae. In *Legumes of the World* (pp. 21-54). Kew, England: Royal Botanic Gardens.
- Shah, A., & Smith, D. L. (2020). Flavonoids in agriculture: Chemistry and roles in biotic and abiotic stress responses, and microbial associations. *Agronomy*, 10(8), 1209. <https://doi.org/10.3390/agronomy10081209>
- Simmonds, M. J. S., Blaney, W. M., Delle Monache, F., & Marini-Bettolo, G. B. (1990). Insect antifeedant activity associated with compounds isolated from species of *Lonchocarpus* and *Tephrosia*. *Journal of Chemical Ecology*, 16(2), 365-380. <https://doi.org/10.1007/BF01020939>
- Sonibare, M. A., Umukoro, S., & Shonibare, E. T. (2012). Antipsychotic property of aqueous and ethanolic extracts of *Lonchocarpus cyanescens* (Schumacher and Thonn.) Benth. (Fabaceae) in rodents. *Journal of Natural Medicines*, 66(1), 127-132. <https://doi.org/10.1007/s11418-011-0562-6>
- Spencer, S. (1996). Developing an understanding of science from the Sierra Leonean traditional Gara dyeing process. *Gender and Science and Technology Association Conference, GASAT 8*, Ahmedabad, January 1996.
- Sule, R. O., Condon, L., & Gomes, A. V. (2022). A common feature of pesticides: Oxidative stress—The role of oxidative stress in pesticide-induced toxicity. *Oxidative Medicine and Cellular Longevity*, 2022, 5563759. <https://doi.org/10.1155/2022/5563759>
- Tuwei, J. K. (2006). Larvicidal and antiparasitoid compounds from *Derris trifoliata*, *Lonchocarpus eriocalyx*, and *Erythrina saculeuxii* (Master's Thesis, University of Nairobi).
- Udeani, G. O., Gerhauser, C. T., Cathy, F. M., Richard, C. K., Jerrome, W., Kinghorn, A. D., Moriarty, R. M., & Pezzuto, J. M. (1997). Cancer chemopreventive activity mediated by deguelin, a naturally occurring rotenoid. *Cancer Research*, 57(16), 3424-3428.
- Vogler, B., Cholewa, L., Schmidt, J., & Setzer, W. (2006). Cytotoxic flavonoids from the bark of *Lonchocarpus haberi* from Monteverde, Costa Rica. *Pharmacologyonline*, 3, 850-855.
- Williams, C. A., & Grayer, R. J. (2004). Anthocyanins and other flavonoids. *Natural Product Reports*, 21, 539-573. <https://doi.org/10.1039/B312631C>
- Wojciechowski, M. F., Lavin, M., & Sanderson, M. J. (2004). A phylogeny of legumes (Leguminosae). *American Journal of Botany*, 91(11), 1846-1862. <https://doi.org/10.3732/ajb.91.11.1846>
- Xueqin, M., Chengjian, Z., Khalid, R., & Luping, Q. (2011). Fabaceae—Traditional uses in Chinese medicine, phytochemistry, and pharmacology. *Journal of Ethnopharmacology*, 138(2), 314-332. <https://doi.org/10.1016/j.jep.2011.09.031>
- Yam-Puc, A., & Peña-Rodríguez, L. M. (2009). Isocordoin derivatives from the root extract of *Lonchocarpus xuul*. *Journal of the Mexican Chemical Society*, 53(1), 12-14.
- Yenesew, A., Induli, M., Derese, S., Midiwo, J. O., Heydenreich, M., Peter, M. G., Akala, H., Wangui, J., Liyala, P., & Waters, N. C. (2004). Antiparasitoid flavonoids from the stem bark of *Erythrina abyssinica*. *Phytochemistry*, 65, 3029-3032. <https://doi.org/10.1016/j.phytochem.2004.06.016>

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2025): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/129910>